



# MVE560 Architectural Geometry, Lecture 1



MVE560, Lecture 1

Mathematical Sciences

**CHALMERS**

6th November 2020

# Outline

Cartesian Coordinates

Some Geometric Primitives

Cylindrical and Spherical Coordinates


# The Cartesian Coordinate System

- The 'usual' coordinate system we use most of the time in  $\mathbb{R}^n$
- Named after French philosopher René Descartes (1596–1650)
- Orthogonal/perpendicular *coordinate axes* — the  $x$ -axis and the  $y$ -axis (and sometimes the  $z$ -axis)
- The *origin* is a special 'reference point' with coordinates  $(0, 0)$  (or  $(0, 0, 0)$  if in 3D)
- May be used for both points and vectors




Image source: Wikipedia

# The Cartesian Coordinate System — Illustration



# Vectors

Vectors live in a *vector space*  $V$  (in our case typically  $V = \mathbb{R}^2$  or  $V = \mathbb{R}^3$ ), equipped with the operations *addition* and *scaling*:



A vector is best thought of as *motion* or a *direction*.

# Vectors in Cartesian Coordinates

In Cartesian coordinates, vector addition and scaling works as follows:

$$(x_1, y_1, z_1) + (x_2, y_2, z_2) = (x_1 + x_2, y_1 + y_2, z_1 + z_2)$$

and

$$\lambda(x, y, z) = (\lambda x, \lambda y, \lambda z).$$

The length, or *norm*, of a vector  $\mathbf{v} = (x, y, z)$  is given by the Pythagorean theorem:

$$\|\mathbf{v}\| = \sqrt{x^2 + y^2 + z^2}.$$

# Points and Vectors

Both vectors and points are often represented using coordinates, e.g.  $(x, y, z)$ , but they are conceptually *very* different!

# Points and Vectors

Both vectors and points are often represented using coordinates, e.g.  $(x, y, z)$ , but they are conceptually *very* different!

- A point has *only* a location, a vector has *no* location



# Points and Vectors

Both vectors and points are often represented using coordinates, e.g.  $(x, y, z)$ , but they are conceptually *very* different!

- A point has *only* a location, a vector has *no* location
- $[\text{vector}] + [\text{vector}] = [\text{vector}]$ , and  $[\text{vector}] - [\text{vector}] = [\text{vector}]$

# Points and Vectors

Both vectors and points are often represented using coordinates, e.g.  $(x, y, z)$ , but they are conceptually *very* different!

- A point has *only* a location, a vector has *no* location
- $[\text{vector}] + [\text{vector}] = [\text{vector}]$ , and  $[\text{vector}] - [\text{vector}] = [\text{vector}]$
- $[\text{point}] - [\text{point}] = [\text{vector}]$ , but  $[\text{point}] + [\text{point}]$  is undefined!

# Points and Vectors

Both vectors and points are often represented using coordinates, e.g.  $(x, y, z)$ , but they are conceptually *very* different!

- A point has *only* a location, a vector has *no* location
- $[\text{vector}] + [\text{vector}] = [\text{vector}]$ , and  $[\text{vector}] - [\text{vector}] = [\text{vector}]$
- $[\text{point}] - [\text{point}] = [\text{vector}]$ , but  $[\text{point}] + [\text{point}]$  is undefined!
- $[\text{point}] + [\text{vector}] = [\text{point}]$

# Points and Vectors

Both vectors and points are often represented using coordinates, e.g.  $(x, y, z)$ , but they are conceptually *very* different!

- A point has *only* a location, a vector has *no* location
- $[\text{vector}] + [\text{vector}] = [\text{vector}]$ , and  $[\text{vector}] - [\text{vector}] = [\text{vector}]$
- $[\text{point}] - [\text{point}] = [\text{vector}]$ , but  $[\text{point}] + [\text{point}]$  is undefined!
- $[\text{point}] + [\text{vector}] = [\text{point}]$
- ...

# Lines

A *line*  $\ell$  consists of all points which can be reached by starting out in a point  $\mathbf{p}_0$  and going in the direction given by a vector  $\mathbf{v} \neq \mathbf{0}$ :

$$\ell : \mathbf{p}(\lambda) = \mathbf{p}_0 + \lambda \mathbf{v}.$$

There exists exactly one line through two distinct points  $\mathbf{p}_1$  and  $\mathbf{p}_2$ :

$$\ell : \mathbf{p}(\lambda) = \mathbf{p}_1 + \lambda(\mathbf{p}_2 - \mathbf{p}_1).$$

# Lines

A *line*  $\ell$  consists of all points which can be reached by starting out in a point  $\mathbf{p}_0$  and going in the direction given by a vector  $\mathbf{v} \neq \mathbf{0}$ :

$$\ell : \mathbf{p}(\lambda) = \mathbf{p}_0 + \lambda \mathbf{v}.$$

There exists exactly one line through two distinct points  $\mathbf{p}_1$  and  $\mathbf{p}_2$ :


$$\ell : \mathbf{p}(\lambda) = \mathbf{p}_1 + \lambda(\mathbf{p}_2 - \mathbf{p}_1).$$

The *line segment* between  $\mathbf{p}_1$  and  $\mathbf{p}_2$  is given by


$$\ell : \mathbf{p}(\lambda) = \mathbf{p}_1 + \lambda(\mathbf{p}_2 - \mathbf{p}_1), \quad 0 \leq \lambda \leq 1.$$

# Triangles (and Other Polygons)


An  $n$ -sided *polygon* is a planar object consisting of *vertices* (corners) which are connected in a particular order by *edges* (line segments):



Triangle



Convex polygon




Non-convex polygon


A polygon is called *convex* if it has no inward 'dents'.

# Triangles (and Other Polygons)


An  $n$ -sided *polygon* is a planar object consisting of *vertices* (corners) which are connected in a particular order by *edges* (line segments):



Triangle



Convex polygon




Non-convex polygon

A polygon is called *convex* if it has no inward 'dents'. Triangles are always convex and planar, and are therefore the polygon most often used to construct things!




# Triangle Meshes



- A *triangle mesh* is a surface consisting of a number of triangles which are joined along their edges.
- With sufficiently many and sufficiently small triangles, triangular meshes can approximate most shapes very well!

# Representing Triangle Meshes



A triangle mesh is often represented using a *vertex list*

$$V = \begin{bmatrix} x_1 & x_2 & \cdots & x_6 \\ y_1 & y_2 & \cdots & y_6 \\ z_1 & z_2 & \cdots & z_6 \end{bmatrix}$$

and a *triangle list*

$$T = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 2 & 4 & 5 & 6 \\ 3 & 5 & 3 & 5 \end{bmatrix},$$

where the indices of the vertices are entered anticlockwise.

# The Platonic Solids




Image source: Wikipedia

- The five *Platonic solids* shown are the only convex solids whose faces are regular polygons
- Many fascinating properties, i.e. symmetries, relations, ...

---

Sutton, *Platonic & Archimedean Solids*, 2002.

# Cylindrical Coordinates

Cylindrical coordinates  $(r, \varphi, z)$  are related to Cartesian coordinates as

$$\begin{cases} x = r \cos \varphi \\ y = r \sin \varphi \\ z = z. \end{cases}$$





Image source: Pottmann *et al.*

# Cylindrical Coordinates

Cylindrical coordinates  $(r, \varphi, z)$  are related to Cartesian coordinates as

$$\begin{cases} x = r \cos \varphi \\ y = r \sin \varphi \\ z = z. \end{cases}$$

Cylindrical coordinates are very useful for describing various kinds of rotational symmetries:







Image source: Pottmann *et al.*

# Spherical Coordinates

Spherical coordinates  $(r, \varphi, \theta)$  are related to Cartesian coordinates as

$$\begin{cases} x = r \cos \varphi \cos \theta \\ y = r \sin \varphi \cos \theta \\ z = r \sin \theta. \end{cases}$$




Image source: Pottmann *et al.*

# Spherical Coordinates

Spherical coordinates  $(r, \varphi, \theta)$  are related to Cartesian coordinates as

$$\begin{cases} x = r \cos \varphi \cos \theta \\ y = r \sin \varphi \cos \theta \\ z = r \sin \theta. \end{cases}$$

Spherical coordinates are useful for 'placing' things in space, e.g. positioning other geometric primitives.




Image source: Pottmann *et al.*