

Chapter 5: FEM for two-point BVP (summary)

November 20, 2020

Goal: We use the theoretical and practical tools from the previous sections to present and analyse various BVP.

- In order to get a FE approximation to the BVP ($a > 0$ and f are given)

$$\begin{cases} -(a(x)u'(x))' = f(x) & \text{for } x \in (0, 1) \\ u(0) = 0 \quad \text{and} \quad u(1) = 0 \end{cases}$$

we proceed as usual:

1. Define the test/trial space $V^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v, v' \in L^2(0, 1), v(0) = v(1) = 0\}$, multiply the DE with a test function $v \in V^0$, integrate over the domain $[0, 1]$ and get the VF

$$\text{Find } u \in V^0 \quad \text{such that} \quad \int_0^1 a(x)u'(x)v'(x) dx = \int_0^1 f(x)v(x) dx \quad \forall v \in V^0.$$

2. Define the finite dimensional space $V_h^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v \text{ is cont. pw. linear on } T_h, v(0) = v(1) = 0\}$, where as usual T_h is a uniform partition with mesh $h = \frac{1}{m+1}$. Observe that $V_h^0 = \text{span}(\varphi_1, \dots, \varphi_m)$ with the hat functions φ_j .

The FE problem then reads

$$\text{Find } U \in V_h^0 \quad \text{such that} \quad \int_0^1 a(x)U'(x)\chi'(x) dx = \int_0^1 f(x)\chi(x) dx \quad \forall \chi \in V_h^0.$$

The above is also called **cG(1) FE** (for linear continuous Galerkin FE).

3. Choosing $\chi = \varphi_i$, writing $U(x) = \sum_{j=1}^m \zeta_j \varphi_j(x)$, and inserting everything into the FE problem gives the following linear system of equations

$$S\zeta = b,$$

where the $m \times m$ **stiffness matrix** S has entries $s_{ij} = \int_0^1 a(x)\varphi_i'(x)\varphi_j'(x) dx$ and the $m \times 1$ **load vector** b has entries $b_i = \int_0^1 f(x)\varphi_i(x) dx$. Formulas for these entries can be found in the book. Solving this system gives the vector ζ and in turns the numerical approximation U .

- The above needs minor adaptations when dealing with other BC. Let us for example derive a FE approximation for the following BVP

$$\begin{cases} -u''(x) + 4u(x) = 0 & \text{for } x \in (0, 1) \\ u(0) = \alpha \quad \text{and} \quad u(1) = \beta, \end{cases}$$

where $\alpha \neq 0$ and $\beta \neq 0$ are given real number. Such boundary conditions are called **non-homogeneous Dirichlet boundary conditions**.

The derivation of a numerical approximation for solutions to the above problem is given by

1. Define the **trial space** $V = \{v: [0, 1] \rightarrow \mathbb{R} : v, v' \in L^2(0, 1), v(0) = \alpha, v(1) = \beta\}$ and the **test space** $V^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v, v' \in L^2(0, 1), v(0) = v(1) = 0\}$. Multiply the DE with a test function $v \in V^0$, integrate over the domain $[0, 1]$ and get the VF

$$\text{Find } u \in V \text{ such that } \int_0^1 u'(x) v'(x) dx + 4 \int_0^1 u(x) v(x) dx = 0 \quad \forall v \in V^0.$$

2. Next, define the finite dimensional spaces

$V_h = \{v: [0, 1] \rightarrow \mathbb{R} : v \text{ is cont. pw. linear on } T_h \text{ and } v(0) = \alpha, v(1) = \beta\}$ and

$V_h^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v \text{ is cont. pw. linear on } T_h, v(0) = v(1) = 0\}$, where as before T_h is a uniform partition with mesh $h = \frac{1}{m+1}$. Observe that $V_h = \text{span}(\varphi_0, \varphi_1, \dots, \varphi_m, \varphi_{m+1}) \subset V$ and $V_h^0 = \text{span}(\varphi_1, \dots, \varphi_m) \subset V^0$ with the hat functions φ_j .

The FE problem then reads

$$\text{Find } U \in V_h \text{ such that } \int_0^1 U'(x) \chi'(x) dx + 4 \int_0^1 U(x) \chi(x) dx = 0 \quad \forall \chi \in V_h^0.$$

3. Choosing $\chi = \varphi_i$, writing $U(x) = \sum_{j=0}^{m+1} \zeta_j \varphi_j(x)$ with $\zeta_0 = \alpha$ and $\zeta_{m+1} = \beta$ (due to the BC), and inserting everything into the FE problem gives the following linear system of equations

$$(S + 4M) \zeta = b,$$

where the $m \times m$ **stiffness matrix** S has entries $s_{ij} = \int_0^1 \varphi_i'(x) \varphi_j'(x) dx$, the $m \times m$ **mass matrix**

M has entries $m_{ij} = \int_0^1 \varphi_i(x) \varphi_j(x) dx$, and the $m \times 1$ **vector** b has entries $b_i = -\alpha(\varphi_0', \varphi_i')_{L^2} - \beta(\varphi_{m+1}', \varphi_i')_{L^2} - 4\alpha(\varphi_0, \varphi_i)_{L^2} - 4\beta(\varphi_{m+1}, \varphi_i)_{L^2}$. Solving this system gives the vector ζ and in turns the numerical approximation U .

- Let us finally consider the problem of finding a numerical approximation of solutions to the BVP

$$\begin{cases} -au''(x) + bu'(x) = r & \text{for } x \in (0, 1) \\ u(0) = 0 \text{ and } u'(1) = \beta, \end{cases}$$

where $\beta \neq 0$, $a, b > 0$, and r are given real number. One has a **homogeneous Dirichlet boundary conditions** for $x = 0$ and **non-homogeneous Neumann boundary conditions** for $x = 1$.

For ease of presentation we take $a = b = r = 1$ and derive a FE approximation as follows

1. Define the space $V = \{v: [0, 1] \rightarrow \mathbb{R} : v, v' \in L^2(0, 1), v(0) = 0\}$. Multiply the DE with a test function $v \in V$, integrate over the domain $[0, 1]$ and get the VF

$$\text{Find } u \in V \text{ such that } (u', v')_{L^2} + (u', v)_{L^2} = \int_0^1 v(x) dx + \beta v(1) \quad \forall v \in V.$$

2. Next, define the finite dimensional space $V_h = \{v: [0, 1] \rightarrow \mathbb{R} : v \text{ is cont. pw. linear on } T_h, v(0) = 0\}$, where as before T_h is a uniform partition with mesh $h = \frac{1}{m+1}$.

Observe that $V_h = \text{span}(\varphi_1, \dots, \varphi_m, \varphi_{m+1}) \subset V$, with the hat functions φ_j .

The FE problem then reads

$$\text{Find } U \in V_h \text{ such that } (U', \chi')_{L^2} + (U', \chi)_{L^2} = \int_0^1 \chi(x) dx + \beta \chi(1) \quad \forall \chi \in V_h.$$

3. Choosing $\chi = \varphi_i$, writing $U(x) = \sum_{j=1}^{m+1} \zeta_j \varphi_j(x)$, observing that φ_{m+1} is a half hat function, and inserting everything into the FE problem gives the following linear system of equations

$$(S + C)\zeta = b,$$

where the $(m+1) \times (m+1)$ **stiffness matrix** S has entries $s_{ij} = \int_0^1 \varphi'_i(x) \varphi'_j(x) dx$, the $(m+1) \times (m+1)$ **convection matrix** C has entries $c_{ij} = \int_0^1 \varphi'_j(x) \varphi_i(x) dx$, and the $(m+1) \times 1$ **vector** b has entries $b_i = \int_0^1 \varphi_i(x) dx + \beta \varphi_i(1)$. Detailed formulas for these entries can be found in the book. Solving this system gives the vector ζ and in turns the numerical approximation U .

- Let $f: (0, 1) \rightarrow \mathbb{R}$ be bounded and continuous. Then, the BVP

$$\begin{cases} -u''(x) = f(x) & \text{for } x \in (0, 1) \\ u(0) = 0 & \text{and } u(1) = 0 \end{cases}$$

is equivalent to the VF

$$\text{Find } u \in \mathcal{C}^2(0, 1) \cap V^0 \text{ such that } (u', v')_{L^2(0, 1)} = (f, v)_{L^2(0, 1)} \text{ for all } v \in V^0.$$

- Poincaré inequality** reads: Let $L > 0$ and consider the open interval $\Omega = (0, L)$. Assume that $u \in H_0^1(\Omega) = \{v: \Omega \rightarrow \mathbb{R} : v, v' \in L^2(\Omega), v(0) = v(L) = 0\}$. Then, one has

$$\|u\|_{L^2(\Omega)} \leq C_L \|u'\|_{L^2(\Omega)}.$$

- A priori error estimate in the energy norm.** Let $f: (0, 1) \rightarrow \mathbb{R}$ be bounded and continuous. Consider the BVP

$$\begin{cases} -u''(x) = f(x) & \text{for } x \in (0, 1) \\ u(0) = 0 & \text{and } u(1) = 0. \end{cases}$$

Denote by U the solution to the corresponding FE problem (cG(1) FE). Assume that $u \in \mathcal{C}^2(0, 1)$. Then, there exists a $C > 0$ such that

$$\|u - U\|_E \leq Ch \|u''\|_{L^2(0, 1)},$$

where $\|v\|_E = \sqrt{(v, v)_E} = \sqrt{(v', v')_{L^2(0, 1)}}$ denotes the energy norm.

Further resources:

- www.simscale.com
- [wiki](#)
- [wiki](#)
- cs.uchicago.edu
- [youtube](#)