

## Chapter 8: The heat equation in 1d (summary)

February 15, 2021

**Goal:** Briefly study the exact solution to some heat equations and present a numerical discretisation.

- Let us start with some **stability estimates** for the following heat equation

$$\begin{cases} u_t(x, t) - u_{xx}(x, t) = f(x, t) & 0 < x < 1, 0 < t \leq T \\ u(0, t) = 0, u_x(1, t) = 0 & 0 < t \leq T \\ u(x, 0) = u_0(x) & 0 < x < 1, \end{cases}$$

where  $u_0$  and  $f$  are given functions.

The solution to the above problem satisfy the following estimates

$$\begin{aligned} \|u(\cdot, t)\|_{L^2(0,1)} &\leq \|u_0\|_{L^2(0,1)} + \int_0^t \|f(\cdot, s)\|_{L^2(0,1)} \, ds. \\ \|u_x(\cdot, t)\|_{L^2(0,1)} &\leq \|u'_0\|_{L^2(0,1)} + \int_0^t \|f(\cdot, s)\|_{L^2(0,1)} \, ds. \end{aligned}$$

When  $f = 0$ , one gets

$$\|u(\cdot, t)\|_{L^2(0,1)} \leq \|u_0\|_{L^2(0,1)} e^{-2t}.$$

When  $f = 0$  and some fixed  $\varepsilon > 0$ , one gets for all  $t \in (0, T]$

$$\int_\varepsilon^t \|u_t(\cdot, s)\|_{L^2(0,1)} \, ds \leq \frac{1}{2} \sqrt{\ln\left(\frac{t}{\varepsilon}\right)} \|u_0\|_{L^2(0,1)}.$$

- Next, we **discretise the inhomogeneous heat equation** with homogeneous Dirichlet boundary conditions

$$\begin{cases} u_t(x, t) - u_{xx}(x, t) = f(x, t) & 0 < x < 1, 0 < t \leq T \\ u(0, t) = u(1, t) = 0 & 0 < t \leq T \\ u(x, 0) = u_0(x) & 0 < x < 1, \end{cases}$$

where  $u_0$  and  $f$  are given functions.

Since it is seldom possible to find the exact solution  $u(x, t)$  to the above problem, we need to find a numerical approximation of it. We proceed as follows

- To get a VF of the heat equation, consider the test/trial space

$V^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v, v' \in L^2(0, 1), v(0) = v(1) = 0\}$ . Then, multiply the DE by a test function  $v \in V^0$ , integrate over  $[0, 1]$ , and use integration by parts to get the VF: For all  $0 < t \leq T$

$$\text{Find } u(\cdot, t) \in V^0 \text{ s.t. } (u_t(\cdot, t), v)_{L^2} + (u_x(\cdot, t), v_x)_{L^2} = (f(\cdot, t), v)_{L^2} \quad \forall v \in V^0 \quad (\text{VF})$$

with the initial condition  $u(x, 0) = u_0(x)$ .

- To get a FE problem, we consider the following subspace of the above space  $V^0$

$V_h^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v \text{ cont. pw. linear on unif. partition } T_h, v(0) = v(1) = 0\} = \text{span}(\varphi_1, \dots, \varphi_m)$ , where  $h = \frac{1}{m+1}$  and  $\varphi_j$  are the hat functions.

The FE problem then reads: For all  $0 < t \leq T$

$$\text{Find } U(\cdot, t) \in V_h^0 \text{ s.t. } (U_t(\cdot, t), \chi)_{L^2} + (U_x(\cdot, t), \chi_x)_{L^2} = (f(\cdot, t), \chi)_{L^2} \quad \forall \chi \in V_h^0 \quad (\text{FE})$$

with the initial condition  $U(x, 0) = \pi_h u_0(x)$  the cont. pw. linear interpolant of  $u_0$ .

3. From the above FE problem, we obtain a system of linear ODE by choosing the test functions  $\chi = \varphi_i$  for  $i = 1, \dots, m$  and writing  $U(x, t) = \sum_{j=1}^m \zeta_j(t) \varphi_j(x)$  with unknown coordinates  $\zeta_j(t)$ .

Inserting everything in (FE), one gets the ODE

$$\begin{aligned} M\dot{\zeta}(t) + S\zeta(t) &= F(t) \\ \zeta(0), \end{aligned} \tag{ODE}$$

where  $M$  is the (already seen)  $m \times m$  mass matrix,  $S$  is the (already seen)  $m \times m$  stiffness matrix,  $F(t)$  is an  $m \times 1$  vector with entries  $F_i(t) = (f(\cdot, t), \varphi_i)_{L^2}$  for  $i = 1, \dots, m$ , the initial condition is given by

$$\zeta(0) = \begin{pmatrix} u_0(x_1) \\ \vdots \\ u_0(x_m) \end{pmatrix},$$

and the unknown vector reads

$$\zeta(t) = \begin{pmatrix} \zeta_1(t) \\ \vdots \\ \zeta_m(t) \end{pmatrix}.$$

4. To find a numerical approximation of  $\zeta(t)$  at some discrete time grid  $t_0 = 0 < t_1 < \dots < t_N = T$ , with  $t_j - t_{j-1} = k = \frac{T}{N}$ , one can for instance use backward Euler scheme which reads

$$\begin{aligned} \zeta^{(0)} &= \zeta(0) \\ (M + kS)\zeta^{(n+1)} &= M\zeta^{(n)} + kF(t_{n+1}) \quad \text{for } n = 0, 1, 2, \dots, N-1. \end{aligned}$$

Solving these linear systems at each time step provides numerical approximations  $\zeta^{(n)} \approx \zeta(t_n)$  that can be inserted in the FE solution to get approximations to the exact solution to the heat equation  $\sum_{j=1}^m \zeta_j(t_n) \varphi_j(x) \approx u(x, t_n)$ .

**Further resources:**

- [wikipedia.org](http://wikipedia.org)
- [math.lamar.edu](http://math.lamar.edu)