

## Chapter 9: The wave equation in 1d (summary)

February 18, 2021

**Goal:** Briefly study the exact solution of the wave equation and present a numerical discretisation of this PDE.

- Consider the (inhomogeneous) **wave equation** with homogeneous Dirichlet boundary conditions

$$\begin{cases} u_{tt}(x, t) - u_{xx}(x, t) = f(x, t) & 0 < x < 1, 0 < t \leq T \\ u(0, t) = u(1, t) = 0 & 0 < t \leq T \\ u(x, 0) = u_0(x) & 0 < x < 1 \\ u_t(x, 0) = v_0(x) & 0 < x < 1, \end{cases}$$

where  $u_0, v_0$  and  $f$  are given (nice) functions.

Introducing a new variable for the velocity  $v = u_t$ , one can rewrite the above wave equation as a system of first order differential equations

$$w_t(x, t) = Aw(x, t) + F(x, t),$$

with  $w(x, t) = \begin{pmatrix} u(x, t) \\ v(x, t) \end{pmatrix}$ ,  $F(x, t) = \begin{pmatrix} 0 \\ f(x, t) \end{pmatrix}$  and the operator  $A = \begin{pmatrix} 0 & 1 \\ \frac{\partial^2}{\partial x^2} & 0 \end{pmatrix}$ .

For the homogeneous wave equation, that is when  $f \equiv 0$  in the above PDE, one has **conservation of the energy**

$$\frac{1}{2} \|u_t(\cdot, t)\|_{L^2}^2 + \frac{1}{2} \|u_x(\cdot, t)\|_{L^2}^2 = \frac{1}{2} \|v_0\|_{L^2}^2 + \frac{1}{2} \|u_0'\|_{L^2}^2 \quad \text{for } 0 \leq t \leq T.$$

- The numerical discretisation of the wave equation is similar to the one for the heat equation:

- The VF reads: Find  $u(\cdot, t) \in V^0$ , for all  $0 < t \leq T$ , such that

$$(u_{tt}(\cdot, t), v)_{L^2} + (u_x(\cdot, t), v_x)_{L^2} = (f(\cdot, t), v)_{L^2}$$

for all test functions  $v \in V^0$  and with initial conditions  $u(x, 0) = u_0(x)$ ,  $u_t(x, 0) = v_0(x)$ .

- The FE problem reads: Find  $U(\cdot, t) \in V_h^0$ , for all  $0 < t \leq T$ , such that

$$(U_{tt}(\cdot, t), \chi)_{L^2} + (U_x(\cdot, t), \chi_x)_{L^2} = (f(\cdot, t), \chi)_{L^2}$$

for all test functions  $\chi \in V_h^0$  and initial conditions  $U(x, 0) = \pi_h u_0(x)$ ,  $U_t(x, 0) = \pi_h v_0(x)$ .

- The linear system of ODEs is given by

$$\begin{aligned} M\dot{\zeta}(t) &= M\eta(t) \\ M\dot{\eta}(t) + S\zeta(t) &= F(t). \end{aligned}$$

Finally, one obtains a numerical approximation of the solution to this ODE by using the Crank–Nicolson scheme with time step  $k$  for instance:

$$\begin{pmatrix} M & -\frac{k}{2}M \\ \frac{k}{2}S & M \end{pmatrix} \begin{pmatrix} \zeta^{(n+1)} \\ \eta^{(n+1)} \end{pmatrix} = \begin{pmatrix} M & \frac{k}{2}M \\ -\frac{k}{2}S & M \end{pmatrix} \begin{pmatrix} \zeta^{(n)} \\ \eta^{(n)} \end{pmatrix} + \begin{pmatrix} 0 \\ \frac{k}{2}(F(t_{n+1}) + F(t_n)) \end{pmatrix}.$$

The Crank–Nicolson scheme preserves a discrete energy (when applied to a homogeneous wave equation).

**Further resources:**

- [wikipedia.org](https://en.wikipedia.org)
- [brilliant.org](https://brilliant.org)
- [math.lamar.edu](https://math.lamar.edu)
- [chem.libretexts.org](https://chem.libretexts.org)