

May 10, 2021

Lecture notes on general and periodic linear ODEs

Plan for the chapter

1. Transition matrix function, existence and equations. Lemma 2.1, p.24, Cor. 2.3, p.26.
2. Grönwall inequality. General case. Lemma. 2.4, p. 27 (skipped)
3. Uniqueness of solutions and dimension of solution space. Th. 2.5, p. 28, Prop. 2.7(1), p.30
4. Group properties of transition matrix function and transition mapping.
(Chapman - Kolmogorov relations) Cor. 2.6, p. 29
5. Fundamental matrix solution and its connection with transition matrix function. Prop. 2.8, p.33
6. Inhomogeneous linear systems. Variation of constant formula (Duhamels formula), general case.
Th. 2.15, p.41, Cor. 2.17.
7. Transition matrix function for periodic linear systems. Formula. 2.31, p. 45.
8. Monodromy matrix and properties of transition matrix function for periodic systems. Th. 2.30, p.
53.
9. Logarithm of a matrix. Prop. 2.29, p. 53.
10. Floquet multipliers and exponents.
11. Boundedness and zero limits for solutions to periodic linear systems. Th. 2.31, p.54. Cor. 2.33, p-
59
12. Existence of periodic solutions to periodic linear systems. Prop. 2.20, p. 45
13. Abel-Liouville's formula. Prop. 2.7(2)., p.30.
14. Conditions for existence of unbounded solutions based on the Abel-Liouville's formula
15. Hill equation and Kapitza pendulum. pp. 55-57.

0.1 Transition matrix function, existence and equations.

The subject of this chapter of lecture notes is general non - autonomous linear systems of ODEs and in particular systems with periodic coefficients and Floquet theory for them.

The general theory for non - autonomous linear systems (linear systems with variable coefficients) is very similar to one for systems with constant coefficients. The existence is established through the solution of the integral form of equations by iterations. Uniqueness is based on a general form of the Grönwall inequality that is also proven here in a very similar fashion. These results lead to the fundamental result on the dimension of the space of solutions that is based on the uniqueness result similarly to the proof for

systems with constant coefficients. The essential difference from the case with constant coefficients is that in the case with variable coefficients one cannot find analytical solutions except some particular cases as systems with triangular matrices.

We consider the I.V.P. in the differential

$$x' = A(t)x(t), \quad x(\tau) = \xi \tag{1}$$

or in the integral form

$$x(t) = \xi + \int_{\tau}^t A(s)x(s)ds \tag{2}$$

with matrix valued function $A : J \rightarrow \mathbb{R}^{N \times N}$ (or $\mathbb{C}^{N \times N}$) that is continuous or piecewise continuous on the interval J .

Here it is important that the initial time τ is an arbitrary real number from J , not just zero.

The solution is defined as a continuous function $x(t)$ on an interval I that includes point $\tau \in I$, acting into \mathbb{R}^N or \mathbb{C}^N , and satisfying the integral equation (2). By a version of Calculus main theorem (Newton-Leibnitz theorem) the solution defined in such a way will satisfy the differential equation (1) in points t where $A(t)$ is continuous.

We remind the following lemma considered in the beginning of the course.

Lemma. The set of solutions \mathcal{S}_{hom} to (2) is a linear vector space.

□

It motivates us to search solution in the form $x(t) = \Phi(t, s)\xi$ where $\Phi(t, s)$ is a continuous matrix valued function on $J \times J$ and ξ is an arbitrary initial data at $t = s : x(s) = \xi$. It implies also that $\Phi(s, s) = I$. Substituting the expression $x(t) = \Phi(t, s)\xi$ into the integral form of the i.V.P.

$$x(t) = \xi + \int_{\tau}^t A(s)x(s)ds$$

we arrive to the vector equation

$$\begin{aligned} \Phi(t, s)\xi &= \xi + \int_s^t A(\sigma)\Phi(\sigma, s)\xi d\sigma \implies \\ \Phi(t, s)\xi &= \left(I + \int_s^t A(\sigma)\Phi(\sigma, s)d\sigma \right) \xi \end{aligned}$$

with arbitrary $\xi \in \mathbb{R}^N$ that implies the matrix equation for $\Phi(t, s)$:

$$\Phi(t, s) = I + \int_s^t A(\sigma)\Phi(\sigma, s)d\sigma \tag{3}$$

or the same equation in differential form valid outside points of discontinuity of $A(t)$:

$$\frac{d}{dt}\Phi(t, s) = A(t)\Phi(t, s); \quad \Phi(s, s) = I.$$

We will solve the equation (3) by means of iterational approximations $M_n(t, s)$ to $\Phi(t, s)$ introduced in the

following way:

$$M_1(t, s) = I; \quad M_{n+1}(t, s) = I + \int_s^t A(\sigma)M_n(\sigma, s)d\sigma, \quad \forall n \in \mathbb{N} \quad (4)$$

Lemma 2.1, p. 24 and **Corollary 2.3**, p. 26 in L&R

For any closed and bounded interval $[a, b] \subset J$ the sequence $\{M_n(t, s)\}$ converges uniformly on $[a, b] \times [a, b]$ to a continuous on $[a, b] \times [a, b]$ matrix valued function $\Phi(t, s)$ that satisfies the integral equation (3).

Point out that outside of points of discontinuity of the matrix $A(t)$ the function $\Phi(t, s)$ satisfies the differential equation $\frac{d}{dt}\Phi(t, s) = A(t)\Phi(t, s)$.

Proof.

The classical idea of the proof is instead of considering $M_n(t, s)$ to consider **telescoping series** with elements $f_{n+1}(t, s) = M_{n+1}(t, s) - M_n(t, s)$, $f_1 = M_1 = I$, with the partial sum that is equal to M_n :

$$M_n = \sum_{k=1}^n f_k$$

where $f_k(t, s)$ is represented as a repeated integral operator from (4):

$$\begin{aligned} M_1(t, s) &= I; \quad M_2(t, s) = I + \int_s^t A(\sigma)M_1(\sigma, s)d\sigma, \\ M_3(t, s) &= I + \int_s^t A(\sigma_1)M_2(\sigma_1, s)d\sigma_1 = \\ &= I + \int_s^t A(\sigma_1) \left[I + \int_s^{\sigma_1} A(\sigma_2)M_1(\sigma_2, s)d\sigma_2 \right] d\sigma_1 \\ &= I + \int_s^t A(\sigma_1)I d\sigma_1 + \int_s^t A(\sigma_1) \int_s^{\sigma_1} A(\sigma_2)M_1(\sigma_2, s)d\sigma_2 d\sigma_1 \\ f_3 &= M_3 - M_2 = \int_s^t A(\sigma_1) \int_s^{\sigma_1} A(\sigma_2)M_1(\sigma_2, s)d\sigma_2 d\sigma_1 \end{aligned}$$

$$f_{n+1}(t, s) = M_{n+1}(t, s) - M_n(t, s) = \int_s^t A(\sigma_1) \int_s^{\sigma_1} A(\sigma_2) \dots \int_s^{\sigma_{n-1}} A(\sigma_n) d\sigma_n \dots d\sigma_2 d\sigma_1$$

for all $(t, s) \in J \times J$, $\forall n \in \mathbb{N}$. Since the matrix $A(t)$ is piecewise continuous on J , it's norm is bounded on any compact subinterval $[a, b] \subset J$:

$$\|A(t)\| \leq K \quad \forall t \in [a, b]$$

We observe using triangle inequality for integrals several times, and the last estimate, that

$$\|f_{n+1}(t, s)\| = \|M_{n+1}(t, s) - M_n(t, s)\| \leq K^n \int_s^t \int_s^{\sigma_1} \dots \int_s^{\sigma_{n-1}} d\sigma_n \dots d\sigma_2 d\sigma_1$$

and after calculating the integral $\int_s^t \int_s^{\sigma_1} \dots \int_s^{\sigma_{n-1}} d\sigma_n \dots d\sigma_2 d\sigma_1 = \frac{1}{n!}(t-s)^n$, based essentially on $\int s^k ds = \frac{s^{k+1}}{k+1}$.

$$\|f_{n+1}(t, s)\| = \|M_{n+1}(t, s) - M_n(t, s)\| \leq \frac{K^n}{n!} |(t-s)|^n \leq \frac{K^n}{n!} (b-a)^n$$

The number series $\sum_{n=0}^{\infty} \frac{K^n}{n!} (b-a)^n$ is convergent to $\exp(K(b-a))$. Therefore by the Weierstrass' criterion the functional series $\sum_{n=1}^{\infty} f_n(t, s)$ converges uniformly on $[a, b] \times [a, b]$ to a limit denoted here by $\Phi(t, s)$. It implies by construction, that the sequence $M_n(t, s)$ converges uniformly on $[a, b] \times [a, b]$ to the limit denoted by $\Phi(t, s)$. Going to the limit in the relation defining iterations (4),

$$\begin{aligned} M_{n+1}(t, s) &= I + \int_s^t A(\sigma) M_n(\sigma, s) d\sigma \implies \\ \Phi(t, s) &= \lim_{n \rightarrow \infty} M_{n+1}(t, s) = I + \int_s^t A(\sigma) \lim_{n \rightarrow \infty} M_n(\sigma, s) d\sigma = I + \int_s^t A(\sigma) \Phi(\sigma, s) d\sigma \end{aligned}$$

we observe that the limit functional matrix $\Phi(t, s)$ satisfies the equation (3). ■

Since the interval $[a, b] \in J$ is arbitrary we may define the function $\Phi : J = J \times J \rightarrow \mathbb{R}^{N \times N}$ (or $\mathbb{C}^{N \times N}$) as the (pointwise) limit:

$$M_n(t, s) \rightarrow \Phi(t, s), \quad n \rightarrow \infty$$

for all $(t, s) \in J \times J$.

Definition. The matrix $\Phi(t, \tau)$ is called **transition matrix function**.

Point out that $\Phi(t, t) = I$. The product $x(t) = \Phi(t, \tau)\xi$ gives by construction the solution to I.V.P. to the equation $x'(t) = A(t)x(t)$ with initial data $x(\tau) = \xi$. In the case when $A(t)$ is only piecewise continuous, $x(t)$ will be continuous and satisfy the corresponding integral equation. It will satisfy the differential equation outside discontinuities of $A(t)$.

Example. For an autonomous linear system with constant matrix A the transition matrix function is $\Phi(t, \tau) = \exp(A(t - \tau))$.

0.2 Grönwall's inequality. Uniqueness of solutions.

Grönwall's lemma. Lemma 2.4., p. 27 in L&R.

(We skip it for now. A simpler version was considered before)

Let $I \subset \mathbb{R}$, be an interval, let $\tau \in I$, and let $g, h : I \rightarrow [0, \infty)$ be continuous nonnegative functions. If for some positive constant $c > 0$,

$$g(t) \leq c + \left| \int_{\tau}^t h(\sigma) g(\sigma) d\sigma \right| \quad \forall t \in I$$

then

$$g(t) \leq c \exp \left(\left| \int_{\tau}^t h(\sigma) d\sigma \right| \right) \quad \forall t \in I$$

Proof.

The proof uses the idea of integrating factor similar to the simpler case with constant $h = \|A\|$ consid-

ered before. Introduce $G, H : I \rightarrow [0, \infty)$ by

$$\begin{aligned} G(t) &= c + \left| \int_{\tau}^t h(\sigma)g(\sigma)d\sigma \right| \\ H(t) &= \left| \int_{\tau}^t h(\sigma)d\sigma \right| \end{aligned}$$

By the hypothesis in the lemma, $0 \leq g(t) \leq G(t)$.

We consider first the case $\tau < t$. Then integrals in the expressions for G and H are nonnegative:

$$G(s) = c + \int_{\tau}^s h(\sigma)g(\sigma)d\sigma; \quad H(s) = \int_{\tau}^s h(\sigma)d\sigma, \quad \forall s \in [\tau, t]$$

Differentiation and the Newton - Leibnitz theorem imply

$$\begin{aligned} G'(s) &= h(s)g(s) \leq h(s)G(s) = H'(s)G(s), \quad \forall s \in [\tau, t] \\ G'(s) - H'(s)G(s) &\leq 0, \quad \forall s \in [\tau, t] \end{aligned}$$

Multiplying the inequality by $\exp(-H(s))$ and observing that

$$(G'(s) - H'(s)G(s)) \exp(-H(s)) = (G(s) \exp(-H(s)))'$$

we arrive to

$$(G(s) \exp(-H(s)))' \leq 0, \quad \forall s \in [\tau, t]$$

Integrating the last inequality from τ to t we arrive to

$$(G(t) \exp(-H(t))) \leq (G(\tau) \exp(-H(\tau))) = c$$

Therefore we arrive to the Grönwalls inequality:

$$(G(t)) \leq c \exp(H(t)) = c \exp\left(\int_{\tau}^t h(\sigma)d\sigma\right)$$

The case when $t < \tau$ is considered similarly by observing that for $t < \tau$

$$G(t) = c + \int_s^{\tau} h(\sigma)(\sigma)d\sigma; \quad H(t) = \int_s^{\tau} h(\sigma)d\sigma, \quad \forall s \in [t, \tau]$$

Do it as an exercise!

Uniqueness of solutions to I.V.P.

Theorem 2.5, p. 28 L&R

Let $(\tau, \xi) \in J \times \mathbb{R}^N (J \times \mathbb{C}^N)$. The function $x(t) = \Phi(t, \tau)\xi$ is a unique solution to the I.V.P. (1). If $y : J_y \rightarrow \mathbb{R}^N$ or (\mathbb{C}^N) is a another solution to (1) $x' = A(t)x(t)$, then $y(t) = x(t)$ for all $t \in J_y$.

Proof.

The fact that $x(t) = \Phi(t, \tau)\xi$ is a solution to I.V:P. follows by construction and from the properties of the transition matrix function. Only uniqueness must be proved. Consider function $e(t) = x(t) - y(t)$ on the interval $J_y \subset J$. By linearity it satisfies the equation

$$e(t) = \int_{\tau}^t A(\sigma)e(\sigma)d\sigma, \quad \forall t \in J_y$$

Applying the triangle inequality for integrals we conclude that

$$\|e(t)\| \leq \int_{\tau}^t \|A(\sigma)\| \|e(\sigma)\| d\sigma, \quad \forall t \in J_y$$

Point out that on an arbitrary bounded closed (compact!) interval $[a, b] \subset J_y$ the piecewise continuous $A(\sigma)$ matrix valued function has a bounded norm $\|A(\sigma)\| < K$. Therefore for any $\tau, t \in [a, b]$

$$\|e(t)\| \leq \|e(\tau)\| + \int_{\tau}^t K \|e(\sigma)\| d\sigma, \quad \forall t, \tau \in [a, b]$$

and by the simple variant of Grönwall 's inequality

$$\|e(t)\| \leq \|e(\tau)\| \exp(K(t - \tau))$$

that we proved before, $\|e(t)\| = 0$ for all $t \in [a, b]$ and therefore $y(t) = x(t)$ for all $t \in J_y$.

0.3 Solution space.

We have considered a particular variant of the following theorem in the case of linear systems of ODEs with constant coefficients. The formulation and the proof we suggested are based only on the fact that the set of solutions \mathbb{S}_h is a linear vector space and on the property of the uniqueness of solutions. We repeat this argument here again with some corollaries about the structure of the transition matrix $\Phi(t, \tau)$.

Proposition 2.7 (1), p.30, L&R.

Let b_1, \dots, b_N be a basis in \mathbb{R}^N (or \mathbb{C}^N) and let $\tau \in J$.

Let $\Phi(t, \tau)$ be a transition matrix to the equation

$$x' = A(t)x$$

with $A(t)$ being a matrix valued function $A : J \rightarrow \mathbb{R}^{N \times N}$ (or $\mathbb{C}^{N \times N}$), piecewise continuous on the interval J .

Then functions $y_j : J \rightarrow \mathbb{R}^N$ (or \mathbb{C}^N) defined as solutions

$$y_j(t) = \Phi(t, \tau)b_j$$

with $j = 1, \dots, N$ to , the equation above form a basis of the solution space \mathbb{S}_h of the equation.

In particular \mathbb{S}_h is N -dimensional and for every solution $x(t) : J \rightarrow \mathbb{R}^N$ (or \mathbb{C}^N) there exist scalars $\gamma_1, \dots, \gamma_N$ such that

$$x(t) = \sum_{j=1}^N \gamma_j y_j(t)$$

for all $t \in J$.

Proof

We can just repeat here the proof that we gave earlier. Point out that it is more general than one given in the course book.

Suppose that at some time t solutions $y_j(t)$ are linearly dependent. It means that there are constants $\{a_j\}_{j=1}^N$ not all zero such that

$$\sum_{j=1}^N a_j y_j(t) = 0$$

at this time. On the other hand there is a solution that satisfies this condition. It is zero solution $x_*(t) = 0$ for all t .

But then these two solutions must coincide because solutions are unique!!! Namely $\sum_{j=1}^N a_j y_j(t) = 0$ for all times including $t = \tau$. Therefore $\sum_{j=1}^N a_j y_j(\tau) = \sum_{j=1}^N a_j b_j = 0$ because b_j are initial conditions at $t = \tau$ for y_j . It is a contradiction because vectors b_j , $j = 1, \dots, N$ are linearly independent. Therefore $y_j(t)$ with $j = 1, \dots, N$ are linearly independent for all t in J . ■

Example.

Calculate the transition matrix function $\Phi(t, s)$ for the system of equations

$$\begin{cases} x'_1 = t x_1 \\ x'_2 = x_1 + t x_2 \end{cases}$$

$$\begin{aligned} x' &= A(t)x; & A(t) &= \begin{bmatrix} t & 0 \\ 1 & t \end{bmatrix} \\ x(\tau) &= \xi \end{aligned}$$

$$x(t) = \Phi(t, \tau)\xi$$

Here the matrix $A(t)$ is triangular.

The system of ODE above has triangular matrix and can be solved recursively starting from the first equation.

The fundamental matrix $\Phi(t, \tau)$ satisfies the same equation, namely

$$\begin{aligned} \frac{d}{dt} \Phi(t, \tau) &= A(t)\Phi(t, \tau) \\ \Phi(\tau, \tau) &= I \end{aligned}$$

$\Phi(t, \tau)$ has columns $\pi_1(t, \tau)$ and $\pi_2(t, \tau)$ that at the time $t = \tau$ have initial values $[1, 0]^T$ and $[0, 1]^T$, because $\Phi(\tau, \tau) = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

We need to find two solutions $\pi_1(t, \tau)$ and $\pi_2(t, \tau)$ that at the time $t = \tau$ have initial values $[1, 0]^T$ and $[0, 1]^T$ to the equation

$$x' = A(t)x;$$

We will use a general solution to the scalar linear equation $x' = p(t)x + g(t)$ with initial data $x(\tau) = x_0$ calculated using the primitive function $\mathbb{P}(t, \tau) = \int_{\tau}^t p(s)ds$ of $p(t)$:

$$x(t) = \exp \{ \mathbb{P}(t, \tau) \} x_0 + \int_{\tau}^t \exp \{ \mathbb{P}(t, s) \} g(s) ds$$

A derivation of this formula using the integrating factor idea follows.

$$\begin{aligned} x' &= p(t)x + g(t), & x_0 &= x(\tau) \\ \mathbb{P}(t, \tau) &= \int_{\tau}^t p(s) ds \\ \exp \{ -\mathbb{P}(t, \tau) \} x' &= \exp \{ -\mathbb{P}(t, \tau) \} p(t)x + \exp \{ -\mathbb{P}(t, \tau) \} g(t) \\ \exp \{ -\mathbb{P}(t, \tau) \} x' - p(t) \exp \{ -\mathbb{P}(t, \tau) \} x &= \exp \{ -\mathbb{P}(t, \tau) \} g(t) \\ \exp \{ -\mathbb{P}(t, \tau) \} x' + (\exp \{ -\mathbb{P}(t, \tau) \})' x &= \exp \{ -\mathbb{P}(t, \tau) \} g(t) \\ [\exp \{ -\mathbb{P}(t, \tau) \} x]' &= \exp \{ -\mathbb{P}(t, \tau) \} g(t) \\ \int_{\tau}^t [\exp \{ -\mathbb{P}(s, \tau) \} x(s)]' ds &= \int_{\tau}^t \exp \{ -\mathbb{P}(s, \tau) \} g(s) ds \\ \exp \{ -\mathbb{P}(t, \tau) \} x(t) - \exp \{ -\mathbb{P}(\tau, \tau) \} x_0 &= \int_{\tau}^t \exp \{ -\mathbb{P}(s, \tau) \} g(s) ds \\ \exp \{ -\mathbb{P}(t, \tau) \} x(t) - \exp \{ 0 \} x_0 &= \int_{\tau}^t \exp \{ -\mathbb{P}(s, \tau) \} g(s) ds \end{aligned}$$

$$\begin{aligned} x(t) &= \exp \{ \mathbb{P}(t, \tau) \} x_0 + \int_{\tau}^t \exp \{ \mathbb{P}(t, \tau) \} \exp \{ -\mathbb{P}(s, \tau) \} g(s) ds \\ x(t) &= \exp \{ \mathbb{P}(t, \tau) \} x_0 + \int_{\tau}^t \exp \{ \mathbb{P}(t, \tau) - \mathbb{P}(s, \tau) \} g(s) ds \\ \mathbb{P}(t, \tau) - \mathbb{P}(s, \tau) &= \int_{\tau}^t p(z) dz - \int_{\tau}^s p(z) dz = \int_{\tau}^t p(z) dz + \int_s^{\tau} p(z) dz = \\ & \int_s^t p(z) dz = \mathbb{P}(t, s) \\ x(t) &= \exp \{ \mathbb{P}(t, \tau) \} x_0 + \int_{\tau}^t \exp \{ \mathbb{P}(t, s) \} g(s) ds; \\ x(\tau) &= x_0 \end{aligned}$$

In the equation

$$x_1' = t x_1$$

the coefficient $p(t) = t$, therefore $\mathbb{P}(t, \tau) = \int_{\tau}^t s ds = \left(\frac{1}{2}s^2\right)\Big|_{\tau}^t = \frac{1}{2}(t^2 - \tau^2)$ and the solution

$$x_1(t) = \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_1(\tau).$$

The second equation

$$x_2' = t x_2 + x_1$$

is similar but inhomogeneous:

$$x_2(t) = \exp(\mathbb{P}(t, \tau))x_2(\tau) + \int_{\tau}^t \exp(\mathbb{P}(t, s))x_1(s)ds.$$

Substituting $\mathbb{P}(t, \tau) = \frac{1}{2}(t^2 - \tau^2)$ we conclude that $\exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_2(\tau) + \int_{\tau}^t \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_1(\tau)ds$

$$\begin{aligned} x_2(t) &= \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_2(\tau) + \int_{\tau}^t \exp\left(\frac{1}{2}(t^2 - s^2)\right) \exp\left(\frac{1}{2}(s^2 - \tau^2)\right)x_1(\tau)ds \\ &= \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_2(\tau) + \int_{\tau}^t \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_1(\tau)ds \end{aligned}$$

And

$$x_2(t) = \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)x_2(\tau) + \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)(t - \tau)x_1(\tau).$$

The fundamental matrix solution $\Phi(t, \tau)$ has columns that are solutions to $x' = A(t)x$ with initial data - that are columns in the unit matrix: $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$,

Taking $x_1(\tau) = 1$ and $x_2(\tau) = 0$ we get $x_1(t) = \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)$ with $x_2(t) = \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)(t - \tau)$

Taking $x_1(\tau) = 0$ and $x_2(\tau) = 1$ we get $x_1(t) = 0$ with $x_2(t) = \exp\left(\frac{1}{2}(t^2 - \tau^2)\right)$ and the fundamental matrix solution in the form

$$\Phi(t, \tau) = \exp\left(\frac{1}{2}(t^2 - \tau^2)\right) \begin{bmatrix} 1 & 0 \\ t - \tau & 1 \end{bmatrix}$$

0.4 Group properties of transition matrix. Chapman - Kolmogorov relations.

remember that in the case with autonomous systems the transition matrix $\Phi(t, \tau) = \exp((t - \tau)A)$.

Therefore in this case


$$\begin{aligned}\Phi(t, \tau) &= \exp \{(t - \tau)A\} = \exp \{(t - \sigma)A\} \exp \{(\sigma - \tau)A\} \\ &= \exp \{(t - \sigma)A + (\sigma - \tau)A\} = \Phi(t, \sigma)\Phi(\sigma, \tau) \\ \Phi(t, \tau) &= \Phi(t, \sigma)\Phi(\sigma, \tau)\end{aligned}$$

The transition matrix $\Phi(t, \tau)$ defines a **transition mapping** $\varphi(t, \tau, \xi)$, that maps initial data ξ at time τ into the state $\varphi(t, \tau, \xi) = x(t) = \Phi(t, \tau)\xi$ of the system at time t .

Let us consider two consecutive solutions $x(t) = \Phi(t, \tau)\xi$ and $y(t) = \Phi(t, \sigma) (\Phi(\sigma, \tau)\xi)$ of the equation $x' = A(t)x(t)$, that continue each other in the time point $t = \sigma$ where the second solution $y(t)$ attains the initial state taken in the point where the first solution $x(t)$ arrives at time $t = \sigma$.

Together with the uniqueness of solutions, this consideration leads to the *group property* of the transition mapping and the transition matrix. The group property means that moving the system governed by the equation $x'(t) = A(t)x(t)$ from time τ to time t is the same as to move it first from time τ to time σ (blue curve) and then to move it without break from time σ to time t (red curve)

$$\Phi(t, \tau)\xi = \Phi(t, \sigma) [\Phi(\sigma, \tau)\xi]$$



Point out that these two "movements" do not need to go both in the positive direction in time as it is in the picture. One of these movements (or both) can go backward in time. Another observation is that the linearity of the system was not essential for this reasoning, only the uniqueness of solutions. We will use a similar argument later for non-linear systems.

We have proven (almost) the following theorem.

Corollary 2.6, p.29, L&R (Chapman - Kolmogorov relations)

For all $t, \sigma, \tau \in J$

$$\Phi(t, \tau) = \Phi(t, \sigma)\Phi(\sigma, \tau), \quad (5)$$

$$\Phi(t, t) = I,$$

$$\Phi(\tau, t)\Phi(t, \tau) = \Phi(\tau, \tau) = I$$

$$\Phi(\tau, t) = (\Phi(t, \tau))^{-1} \quad (6)$$

Proof.

The first statement has been proven already. The second follows from the integral equation for the transition matrix. The third one follows from the first two. We apply the first statement $\Phi(t, \tau)\Phi(\tau, t) = \Phi(t, t) = I$ therefore $\Phi(\tau, t)$ is the right inverse of $\Phi(t, \tau)$. The same argument for this expression with t and τ changed their roles leads to that $\Phi(\tau, t)$ is the left inverse of $\Phi(t, \tau)$. ■

0.5 Fundamental matrix solution.

Introducing the transition matrix function $\Phi(t, \tau)$ for non-autonomous system of equations was similar to considering $\exp(A(t - \tau))$ for autonomous linear systems. We have got a solution to an arbitrary I.V.P. by multiplying arbitrary initial data $x(\tau) = \xi$ with the the transition matrix function: $x(t) = \Phi(t, \tau)\xi$.

On the other hand we could construct a general solution to an autonomous linear system just by taking a linear combination of N linearly independent solutions to the system, because the dimension of the solution space is equal to N .

The situation is exactly the same for non-autonomous linear systems with the difference that we in general cannot find a basis for the space of solutions analytically. It is possible only in some particular cases, for example for a triangular matrix $A(t)$.

Definition.

The functon $t \mapsto \Psi(t) \in \mathbb{R}^{n \times n}$ is called the **fundamental matrix solution** for the system $x' = A(t)x$, $x \in \mathbb{R}^n$ if it's columns $\Psi_k(t)$, $k = 1, \dots, N$ are linearly independent solutions to the system (and therefore build a basis to the solution space): $\Psi'_k(t) = A(t)\Psi_k(t)$

It follows from the definition of the matrix product that

$$\Psi'(t) = A(t)\Psi(t)$$

General solution to the system is a linear combination of these vector valued functions:

$$x(t) = \Psi(t)C$$

with an arbitrary constant vector $C \in \mathbb{R}^N$.

The fundamental matrix solution $\Psi(t)$ is an invertible matrix for all t because it's columns are linearly independent for all t .

There is a simple connection between an arbitrary fundamental matrix solution $\Psi(t)$ and the transition matrix $\Phi(t, \tau)$.

Proposition 2.8 , p. 33

$$\Phi(t, \tau) = \Psi(t)\Psi^{-1}(\tau)$$

Proof.

The product $X(t, \tau) = \Psi(t)\Psi^{-1}(\tau)$ satisfies the equation

$$X'(t, \tau) = A(t)X(t, \tau)$$

in all points $t \in J$ where $A(t)$ is continuous, because each column in $\Psi(t)$ does it. On the other hand $\Psi(\tau)\Psi^{-1}(\tau) = I$. Therefore $X(t, \tau) = \Psi(t)\Psi^{-1}(\tau)$ satisfies the integral equation

$$X(t, \tau) = I + \int_{\tau}^t A(\sigma)X(\sigma, \tau)d\sigma$$

in all points $t \in J$ because each column in $\Psi(t)$ does it. The same equations are satisfied by $\Phi(t, \tau)$. By the uniqueness of solutions to linear systems of ODEs $\Phi(t, \tau) = X(t, \tau) = \Psi(t)\Psi^{-1}(\tau)$.

This proposition shows an alternative way to calculate the transition matrix solution, because sometimes it is easier to find some basis for the space of solutions and to put it into a matrix $\Psi(t)$ instead of solving the matrix equation for $\Phi(t, \tau)$.

0.6 Abel - Liouville's formula.

Lemma about the derivative of a determinant of a matrix valued function.

Let $B : J \rightarrow \mathbb{R}^{N \times N}$ be differentiable. Then the derivative of it's determinant satisfies the following formula

$$(\det(B(t)))' = \sum_{k=1}^N \det(U_k(B))$$

where matrices $U_k(B)$ have the same columns $b_k(t)$ as the matrix $B(t) = [b_1(t), \dots, b_N(t)]$ except the k -th column exchanged by the column of derivatives of the k -th column in $B(t)$.

$$U_k(B) = \left[b_1(t), \dots, \left[\frac{d}{dt} b_k(t) \right], \dots, b_N(t) \right]$$

A similar relation can be written for rows instead of columns.

An elementary proof can be carried out using the definition of derivative as a limit of a finite difference and repeated application of the addition formula for determinants. **Prove it as an exercise on properties of determinants!**

Consider a homogeneous linear system of ODEs $x'(t) = A(t)x(t)$ and N solutions $y_1(t), y_2(t), \dots, y_N(t)$

to it. Consider the matrix $Y(t)$ having these solutions as it's columns:

$$Y(t) = [y_1(t), y_2(t), \dots, y_N(t)]$$

Definition.

The determinant

$$w(t) = \det Y(t) = \det [y_1(t), y_2(t), \dots, y_N(t)]$$

is called **Wronskian** associated with solutions $y_1(t), y_2(t), \dots, y_N(t)$.

Proposition 2.7 part (2) - Abel - Liouville's formula

Wronskian $w(t)$ associated with solutions $y_1(t), y_2(t), \dots, y_N(t)$ to the system $x'(t) = A(t)x(t)$ satisfies the following relations:

$$w(t) = w(\tau) \det \Phi(t, \tau)$$

In points t where $A(t)$ is continuous it satisfies the differential equation

$$w'(t) = \text{tr}(A(t))w(t)$$

and therefore with initial value for $w(\tau)$ at time τ :

$$w(t) = w(\tau) \exp \left(\int_{\tau}^t \text{tr}(A(s)) ds \right) \tag{7}$$

for all $t \in J$. \square

Proof.

We use here that $y_k(t) = \Phi(t, \tau)y_k(\tau)$ and therefore $Y(t) = \Phi(t, \tau)Y(\tau)$. It implies that

$$w(t) = \det Y(t) = \det (\Phi(t, \tau)Y(\tau)) = \det Y(\tau) \det \Phi(t, \tau) = w(\tau) \det \Phi(t, \tau)$$

giving the first statement of the Proposition.

We denote by $\varphi_k(t)$ columns in $\Phi(t, \tau)$, so that $\Phi(t, \tau) = [\varphi_1(t), \varphi_2(t), \dots, \varphi_N(t)]$. Then we apply the **Lemma about the derivative of a determinant of a matrix valued function** to the case $B(t) = \Phi(t, \tau)$. A direct substitution implies that

$$\frac{\partial}{\partial t} (\det \Phi(t, \tau)) = \sum_{k=1}^N \det (U_k(\Phi(t, \tau))) = \sum_{k=1}^N \det \left(\left[\varphi_1(t), \dots, \frac{\partial}{\partial t} (\varphi_k(t)), \dots, \varphi_N(t) \right] \right)$$

where the k -th column in $U_k(\Phi(t, \tau))$ is $\frac{\partial}{\partial t} (\varphi_k(t))$ and other columns are columns $\varphi_j(t)$, $j \neq k$, $j = 1, \dots, N$ from $\Phi(t, \tau)$.

$\frac{\partial}{\partial t} (\varphi_k(t)) = A(t)\varphi_k(t)$, because $\varphi_k(t)$ are solutions to the system $x'(t) = A(t)x(t)$. We assume here that τ is not a point of discontinuity for $A(t)$. It leads to the more explicit expression:

$$\frac{\partial}{\partial t} (\det \Phi(t, \tau)) = \sum_{k=1}^N \det (U_k(\Phi(t, \tau))) = \sum_{k=1}^N \det ([\varphi_1(t), \dots, A(t) (\varphi_k(t)), \dots, \varphi_N(t)])$$

Setting $t = \tau$, into the last formula for we arrive to

$$\left. \frac{\partial}{\partial t} (\det \Phi(t, \tau)) \right|_{t=\tau} = \sum_{k=1}^N \det ([e_1, \dots, A(\tau)e_k, \dots, e_N])$$

because $\Phi(\tau, \tau) = I = [e_1, \dots, e_k, \dots, e_N]$. Observe that $A(\tau)e_k = [A(\tau)]_k$ - is the k -th column in $A(\tau)$. Matrices under the determinant sign in the last formula are diagonal with all elements equal to one, except the k -th kolumn equal to $[A(\tau)]_k$. See the expression for this matrix in the case of $N = 5, k = 4$.

$$\begin{bmatrix} 1 & 0 & 0 & A_{14} & 0 \\ 0 & 1 & 0 & A_{24} & 0 \\ 0 & 0 & 1 & A_{34} & 0 \\ 0 & 0 & 0 & A_{44} & 0 \\ 0 & 0 & 0 & A_{54} & 1 \end{bmatrix}$$

Applying the standard expansion formula along the first column of the determinant we conclude that the determinants in the right hand side of the formula for $\left. \frac{\partial}{\partial t} (\det \Phi(t, \tau)) \right|_{t=\tau}$ are equal to the product of diagonal elements:

$$\det ([e_1, \dots, A(\tau)e_k, \dots, e_N]) = A(\tau)_{kk}$$

For example for $N = 5, k = 3$.

$$\begin{aligned} \det [e_1, \dots, [A(\tau)]_k, \dots, e_N] &= \det \begin{bmatrix} 1 & 0 & A_{13} & 0 & 0 \\ 0 & 1 & A_{23} & 0 & 0 \\ 0 & 0 & A_{33} & 0 & 0 \\ 0 & 0 & A_{43} & 1 & 0 \\ 0 & 0 & A_{53} & 0 & 1 \end{bmatrix}, \quad k = 3 \\ &= 1 \times 1 \times A_{33} \times 1 \times 1 = A_{33} \end{aligned}$$

Therefore

$$\left. \frac{\partial}{\partial t} (\det \Phi(t, \tau)) \right|_{t=\tau} = \sum_{k=1}^N \det ([e_1, \dots, [A(\tau)]_k, \dots, e_N]) = \sum_{k=1}^N A_{kk}(\tau) = \text{tr} A(\tau)$$

Therefore we derived the differential equation for wronskian for arbitrary:

$$w'(\tau) = w(\tau)\text{tr}A(\tau)$$

The argument given here applies to any $\tau \in J$ that is not a point of discontinuity for $A(t)$. The expression

$$\begin{aligned} w(t) &= w(\tau) \exp \left(\int_{\tau}^t \operatorname{tr}(A(s)) ds \right) \\ w(t) &= \det Y(t) \end{aligned}$$

follows by integration of the differential equation for $w(t)$ using the method of integrating factor applied to a scalar first order linear equation. In the case when $A(\dots)$ has finite number of discontinuities at the interval (τ, t) the formula follows by applying the same argument consequently on each interval of continuity. ■

Interesting observations with application of Abel - Liouville's formula.

The geometric meaning of determinant $\det(C)$ of the matrix $C = [c_1, \dots, c_N]$ with columns c_1, \dots, c_N is volume of the parallelepiped V build on vectors c_1, \dots, c_N :

$$|\det(C)| = \operatorname{vol}(V)$$

One can define V formally as $V = \left\{ x \in \mathbb{R}^N : x = \sum_{k=1}^N a_k c_k, a_k \in [0, 1], k = 1, \dots, n \right\}$.

It implies that the Abel - Liouville's formula gives an exact description of how for example the volume of a unique cube build on standard basis vectors e_1, \dots, e_N given at the initial time τ is changing by the "flow" described by the transition matrix function $\Phi(t, \tau)$.

Lecture 11 Summary of Lecture 10 on non-autonomous linear systems

1. Transition matrix function $\Phi(t, \tau)$ representing a solution to a non-autonomous system $x'(t) = A(t)x(t)$, $x(\tau) = \xi \implies x(t) = \Phi(t, \tau)\xi$.

2. This matrix defines a linear **transition mapping** $\varphi(t, \tau, \xi)$, that maps initial data ξ at time τ into the state $\varphi(t, \tau, \xi) = x(t) = \Phi(t, \tau)\xi$ of the system at time t .

3. Space \mathbb{S}_h of solutions to $x'(t) = A(t)x(t)$.

4. Uniqueness of solutions to I.V.P (by Grönwall's inequality)

5. Construction of the basis to the solution space \mathbb{S}_h .

6. Fundamental matrix solution $\Psi(t, \tau)$ and it's connection with transition matrix function $\Phi(t, \tau)$.

Using fundamental matrix solution for constructing general solution.

7. Chapman-Kolmogorov relations: $\Phi(t, \tau) = \Phi(t, \sigma)\Phi(\sigma, \tau)$

8. Wronskian $w(t) = \det [y_1(t), y_2(t), \dots, y_N(t)]$ of a set of solutions $[y_1(t), y_2(t), \dots, y_N(t)]$ and the Abel-Liouville's formulas:

$$\begin{aligned} w'(t) &= \operatorname{tr}(A(t))w(t) \\ w(t) &= w(\tau) \exp \left(\int_{\tau}^t \operatorname{tr}(A(s)) ds \right) \end{aligned}$$

8. Geometric meaning of Abel- Liouville's formulas.

0.7 Non-homogeneous linear systems and Duhamel's formula in the general

case.

We consider the I.V.P. for non-homogeneous linear system

$$x'(t) = A(t)x(t) + b(t), \quad x(\tau) = \xi, \quad (\tau, \xi) \in J \times \mathbb{R}^N (J \times \mathbb{C}^N)$$

We suppose here that $A : J \rightarrow \mathbb{R}^{N \times N}$ (or $\mathbb{C}^{N \times N}$) is continuous or piecewise continuous and denote by $\Phi(t, \tau)$ the transition matrix function generated by $A(t)$. We rewrite the I.V.P. for the system also in integral form

$$x(t) = \xi + \int_{\tau}^t (A(\sigma)x(\sigma) + b(\sigma)) d\sigma,$$

that allows to consider continuous solutions in the case when A is only peacewise continuous. In this case solutions satisfy the differential form of the problem in between of discontinuities of A .

Theorem 2.15, p. 41 L&R

Let $(\tau, \xi) \in J \times \mathbb{R}^N$. The function

$$x(t) = \Phi(t, \tau)\xi + \int_{\tau}^t \Phi(t, \sigma)b(\sigma)d\sigma,$$

is a unique solution to the I.V.P. above.

Proof. A simpler proof can be given for points t outside the discontinuities of A . Then the result for an arbitrary time interval follows by repeated argument for each subinterval $[\tau, t]$ where A is continuous.

Apply the Chapman-Kolmogorov relation to the transition matrix under the integral: $\Phi(t, \sigma) = \Phi(t, 0)\Phi(0, \sigma)$ and calculate derivative of the integral in the expression for the solution.

$$\begin{aligned} & \frac{d}{dt} \left(\int_{\tau}^t \Phi(t, \sigma)b(\sigma)d\sigma \right) \\ &= \frac{d}{dt} \left(\int_{\tau}^t \Phi(t, 0)\Phi(0, \sigma)b(\sigma)d\sigma \right) = \frac{d}{dt} \left(\Phi(t, 0) \int_{\tau}^t \Phi(0, \sigma)b(\sigma)d\sigma \right) \\ &= \left(\frac{d}{dt} \Phi(t, 0) \right) \int_{\tau}^t \Phi(0, \sigma)b(\sigma)d\sigma + \left(\Phi(t, 0) \frac{d}{dt} \left(\int_{\tau}^t \Phi(0, \sigma)b(\sigma)d\sigma \right) \right) \\ &= A\Phi(t, 0) \int_{\tau}^t \Phi(0, \sigma)b(\sigma)d\sigma + \Phi(t, 0)\Phi(0, t)b(t) \end{aligned}$$

Observe that by Chapman -Kolmogorov relation $\Phi(t, 0)\Phi(0, t) = \Phi(t, t) = I$, and $\Phi(t, 0)\Phi(0, \sigma) = \Phi(t, \sigma)$. It implies simplifications in the last formula and finally

$$\frac{d}{dt} \left(\int_{\tau}^t \Phi(t, \sigma)b(\sigma)d\sigma \right) = A \left(\int_{\tau}^t \Phi(t, \sigma)b(\sigma)d\sigma \right) + b(t)$$

Therefore $\int_{\tau}^t \Phi(t, \sigma)b(\sigma)d\sigma$ is the solution to the inhomogeneous equation with initial condition zero. Adding the solution $\Phi(t, \tau)\xi$ to the homogeneous equation, satisfying the initial condition $\Phi(\tau, \tau)\xi = \xi$ we

conclude that $x(t) = \Phi(t, \tau)\xi + \int_{\tau}^t \Phi(t, \sigma)b(\sigma)d\sigma$, is a solution to the I.V.P. above. The uniqueness follows if we consider difference between two solutions $x(t)$ and $y(t)$ with the same initial condition: $z(t) = x(t) - y(t)$ that evidently satisfies the homogeneous equation $z'(t) = A(t)z(t)$ and the zero initial condition $z(\tau) = 0$. The known result for homogeneous linear systems based on Grönwall's inequality implies that $z(t) = 0$ on J .

Another proof based on the integral formulation of the problem and on the explicit checking using Fubini theorem that $x(t)$ expressed as in the formulation of the theorem satisfies it, is given in the book on the page 41.

1 Systems with periodic coefficients: Floquet theory

We consider here linear homogeneous systems of ODE's with $J = \mathbb{R}$ and a continuous or piecewise continuous matrix $A : \mathbb{R} \rightarrow \mathbb{R}^{N \times N}$ (or $\mathbb{C}^{N \times N}$), with period $p > 0$:

$$x'(t) = A(t)x(t), \quad A(t+p) = A(t), \quad \forall t \in \mathbb{R}$$

Let Φ be a transition matrix generated by a periodic $A(t)$.

Shifting invariance property.(formula 2.31, p. 45 in L.R.)

We are going to prove an important *shifting invariance property* of this transition matrix function, namely that

$$\Phi(t+p, \tau+p) = \Phi(t, \tau) \tag{8}$$

Structure of the transition matrix for a time interval including a finite number of periods.(formula 2.32, p. 45 in L.R.)

(Motivation to introducing the monodromy matrix)

Another property specifying further how the periodicity of the system influences properties of solutions.

$$\Phi(t+p, \tau) = \Phi(t, 0) [\Phi(p, 0)] \Phi(0, \tau) \tag{9}$$

$$\Phi(t+np, \tau) = \Phi(t, 0) [\Phi(p, 0)]^n \Phi(0, \tau) \tag{10}$$

for any $(t, \tau) \in \mathbb{R} \times \mathbb{R}$.

Definition of the Monodromy matrix

The transition matrix $\Phi(p, 0)$ for a periodic linear system with period p is called the **monodromy matrix** (this standard notion is not used in the course book)

Proofs

Proof of the shifting invariance property.

This first property becomes untuitively clear after followig arguments:

$$\begin{aligned} & \Phi(t + np, \tau) \stackrel{Ch.-Kol.}{=} \Phi(t + np, \tau + np) \Phi(\tau + np, \tau) \stackrel{Shift}{=} \Phi(t, \tau) \Phi(\tau, \tau - np) \\ & \stackrel{Ch.-Kol.}{=} \Phi(t, \tau) \Phi(\tau, 0) \Phi(0, \tau - np) \stackrel{Ch.-Kol.}{=} \Phi(t, 0) \Phi(np, \tau) \\ & \stackrel{Ch.-Kol.}{=} \Phi(t, 0) \Phi(np, 0) \Phi(0, \tau) \end{aligned}$$

and from the observation that $\Phi(np, 0) \stackrel{Ch.-Kol.}{=} \Phi(np, np-p) \dots \Phi(kp, kp-p) \dots \Phi(2p, p) \Phi(p, 0) \stackrel{shifting_for_each}{=} [\Phi(p, 0)]^n$ that follows from the Chapman-Kolmogorov relation and from the fact that $\Phi(t, 0)$ satisfies the same equation on each interval $[kp, (k+1)p]$, (shift invariance property) because $A(t) = A(t+p)$ is a periodic matrix with period p . It implies the desired formula:

$$\Phi(t + np, \tau) = \Phi(t, 0) [\Phi(p, 0)]^n \Phi(0, \tau)$$

■

Example illustrating ideas of Floquet theory on a scalar linear equation.

Consider the following scalar linear equation with periodic coefficient $A(t) = (\sin(4t) - 0.1)$ with period $p = 0.5\pi$:

$$\frac{dx}{dt} = (\sin(4t) - 0.1) x,$$

We will find the monodromy matrix for this simple equation and demonstrate all objects related to the Floquet theorem that follows.

The exact general solution is:

$$x(t) = C \exp(-0.25 \cos(4t) - 0.1t)$$

with arbitrary constant C , can be found by the method with integrating factor.

$-0.25 \cos(4t) - 0.1t$ is the primitive function of the coefficient $(\sin(4t) - 0.1)$ in front of x in the equation.

To find the solution equal to 1 at $t = 0$ that is the transition "matrix" in the scalar case, we calculate the expression $\exp(-0.25 \cos(4.0t)) e^{-0.1t} \Big|_{t=0} = 0.7788$ and choose $C = \frac{1}{0.7788}$ in the expression for the general solution $x(t)$.

The transition "matrix" is:

$$\Phi(t, 0) = \frac{1}{0.7788} \exp(-0.25 \cos(4.0t)) e^{-0.1t}$$

The period of the coefficient in the system is $p = 0.5\pi$ and the **monodromy matrix** is $\Phi(p, 0) = \Phi(0.5\pi, 0)$:

$$\Phi(p, 0) = \frac{1}{0.7788} \exp(-0.25 \cos(4.0t)) e^{-0.1t} \Big|_{t=0.5\pi} = 0.85464$$

The eigenvalue μ of the (1x1) "monodromy matrix" $\Phi(p, 0)$ coincides with its value: $\mu = 0.85464 < 1$ and is strictly less than 1.

Consider the logarithm $G = \ln(\Phi(p, 0))$ of the **monodromy matrix** $\Phi(p, 0)$:

$$G = \ln(\Phi(p, 0)) = \ln\left(\frac{1}{0.7788} \exp(-0.25 \cos(4.0t)) e^{-0.1t}\right) \Big|_{t=0.5\pi} = -0.15708$$

$$F = \frac{G}{p} = \frac{-0.15708}{0.5\pi} = -0.1 < 0$$

Therefore the eigenvalue $\lambda = -0.1$ of the "matrix"

$$F = \frac{1}{p}G = \frac{1}{p} \ln(\Phi(p, 0))$$

is negative.

The transition matrix to the linear homogeneous "system"

$$y'(t) = Fy(t)$$

is

$$\exp(Ft) = \exp\left(t \frac{G}{p}\right) = \exp(-0.1t).$$

Compare black and orange graphs for $\exp\left(t \frac{G}{p}\right)$ and for $\Phi(t, 0) = \frac{1}{0.7788} \exp(-0.25 \cos(4.0t)) e^{-0.1t}$. Observe that $\exp\left(t \frac{G}{p}\right)$ and $\Phi(t, 0)$ coincide in points $t = pn = (0.5\pi)n$, $n = 1, 2, 3, \dots$


Introduce a "corrector" multiplier $\Theta(t)$ introduced so that

$$\Phi(t, 0) = \Theta(t) \exp\left(t \frac{G}{p}\right)$$

Observe that

$$\Theta(t) = \frac{1}{0.7788} \exp(-0.25 \cos(4.0t))$$

is a $p = 0.5\pi$ - periodic function equal to 1 in all points $t = pn = (0.5\pi)n$, $n = 1, 2, 3, \dots$ (red graf).



We are going to observe soon that a similar representation of the transition matrix $\Phi(t, 0)$ is possible for the transition matrix $\Phi(t, 0)$ of an arbitrary periodic linear systems of ODEs.

The main idea of the Floquet theory.

The monodromy matrix $\Phi(p, 0)$ is a particular transition matrix that maps initial data ξ at time $\tau = 0$ to the state of the system $x(p)$ after one period p . A particular property of this matrix in the case of periodic systems is that similar mapping to the state $x(t)$ at the time $t = np$ equal to n periods p is just

$$\Phi(n \cdot p, 0) = [\Phi(p, 0)]^n$$

This property is similar to properties of autonomous linear systems where $\Phi(t, 0) = \exp(At)$ and therefore

$$\Phi(n \cdot p, 0) = \exp(A(n \cdot p)) = [\exp(A(p))]^n = [\Phi(p, 0)]^n \tag{11}$$

that follows from the factorisation property of the exponent of two commuting matrices:

$$\exp(A + B) = \exp(A) \exp(B)$$

In the case of periodic systems this factorisation applies only for shifts in time that are integer numbers of periods. But it is still a remarkable property. The behaviour of solutions is described by a repeated multiplication by a constant matrix in certain time points: $p, 2p, 3p, \dots$:

$$x'(t) = A(t)x(t), \quad x(0) = \xi.$$

$$x(np) = [\Phi(p, 0)]^n \xi, \quad n = 0, 1, 2, \dots$$

The first idea of the Floquet theory is to represent $x(np)$ at times $t = np$ similarly as for autonomous systems, namely with the help of an exponent of some (unknown at the moment) constant matrix F times the time argument: $t = np$.

$$x(np) = [\Phi(p, 0)]^n \xi = \exp(np F)\xi = [\exp(p F)]^n \xi$$

It means that the matrix F in such representation must satisfy the relation

$$\Phi(p, 0) = \exp(pF).$$

Therefore the matrix pF must be something like the logarithm of the monodromy matrix:

$$pF = \log(\Phi(p, 0))$$

Definition. A matrix $G \in \mathbb{C}^{N \times N}$ is called a **logarithm of the matrix** $H \in \mathbb{C}^{N \times N}$ if

$$H = \exp(G)$$

We write in this case $G = \log(H)$.

We are going to prove soon that for any non-singular matrix H there is a logarithm $\log(H)$ in this sense. Point out that the monodromy matrix $\Phi(p, 0)$ is always non-singular, because columns in a transition matrix $\Phi(t, 0)$ are always linearly independent (check yourself: why?)

The logarithm of a matrix is not uniquely defined in the same way as it is not unique for complex and real numbers z :

$$\ln(z) = \ln(|z|) + i \arg(z) \tag{12}$$

because the argument $\arg(z)$ of a complex number is defined only up to $2\pi k$, $k = \pm 1, \pm 2, \dots$

One can choose a unique branch for the logarithm function, called the *principle logarithm* or $\text{Log}(z)$ by choosing the argument in the last formula (12) for example only in the interval $(-\pi, \pi]$.

We will suspend the discussion of matrix logarithm now and will consider first an application of it to the analysis of solutions to periodic linear systems of ODEs.

The main idea in the Floquet theory is the "approximation" of the transition matrix $\Phi(t, 0)$ for a periodic linear system with matrix $A(t) = A(p + t)$ by the transition matrix $\exp(tF)$ for an autonomous system

$$y'(t) = [F]y(t)$$

with the constant matrix $F = \left[\frac{1}{p}G \right]$ where

$$G = \log(\Phi(p, 0)) \tag{13}$$

$$\exp(G) = \Phi(p, 0) \tag{14}$$

$$\exp(pF) = \Phi(p, 0) \tag{15}$$

$$\exp((np)F) = [\Phi(p, 0)]^n = \Phi(np, 0) \tag{16}$$

$\Phi(p, 0)$ is considered as a transition matrix for an autonomous system.

These two transition matrices coincide in points $t = 0, p, 2p, 3p, \dots$

$$\Phi(np, 0) = [\Phi(p, 0)]^n = \exp((np)[F]) \tag{17}$$

The "deviation" of $\Phi(t, 0)$ from $\exp(tF)$ in intermediate points within one period can be expressed by a factor $\Theta(t)$ so that

$$\Phi(t, 0) = \Theta(t) \exp(tF)$$

$$\Theta(t) = \Phi(t, 0) \exp(-tF)$$

The matrix function $\Theta(t)$ must be equal to the unit matrix I in the points $t = 0, p, 2p, \dots$ because in these points these two transition functions coincide by construction and $\exp(-tF)$ is inversion of $\Phi(t, 0)$, see (17).

The exact formulation of the properties of such factorization is given in the following Theorem by Floquet.

Lecture 12

Summary of Lecture 11

on periodic systems and Floquet theorem

1. Shifting property of transition matrix for periodic systems:


$$\Phi(t + p, \tau + p) = \Phi(t, \tau)$$

2. Structure formula for transition matrix of periodic systems

$$\Phi(t + p, \tau) = \Phi(t, 0) [\Phi(p, 0)] \Phi(0, \tau) \tag{18}$$

$$\Phi(t + np, \tau) = \Phi(t, 0) [\Phi(p, 0)]^n \Phi(0, \tau) \tag{19}$$

3. Picture of the transition mapping $\varphi(t, \tau, \xi) = \Phi(t, \tau)\xi$ generated by the transition matrix $\Phi(t, \tau)$ to a differential equation $x'(t) = A(t)x(t)$, with periodic coefficients $A(t + p) = A(t)$



Theorem 2.30 , p. 53. Floquet theorem

Consider a periodic system $x'(t) = A(t)x(t)$, with period p : $A(t) = A(t + p)$

Let $G \in \mathbb{C}^{N \times N}$ be a logarithm of the monodromy matrix $\Phi(p, 0)$.

$$G = \log(\Phi(p, 0))$$

There exists a periodic with period p piecewise continuously differentiable function $\Theta(t) : \mathbb{R} \rightarrow \mathbb{C}^{N \times N}$, with $\Theta(0) = I$ and $\Theta(t)$ non-singular (invertible, all eigenvalues are non-zero) for all t , such that

$$\Phi(t, 0) = \Theta(t) \exp\left(\frac{t}{p}G\right), \quad \forall t \in \mathbb{R} \tag{20}$$

Proof.

We remind the main property (9) of the monodromy matrix for $\tau = 0$:

$$\Phi(t + p, 0) = \Phi(t + p, p)\Phi(p, 0) = \Phi(t, 0)\Phi(p, 0)$$

where we applied first the Chapman Kolmogorov relation (5) and then the shift invariance (8) of the transition matrix function $\Phi(t, \tau)$ for a periodic linear system.

We denote $\frac{1}{p}G$ by F for convenience, so that $G = pF$, and define the matrix function $\Theta(t)$ after the desired relation (20)

$$\Theta(t) = \Phi(t, 0) \exp\left(-\frac{t}{p}G\right) = \Phi(t, 0) \exp(-tF)$$

The matrix function $\Theta(t)$ is well defined in such a way. The problem is to show that it has desired properties: p -periodicity and satisfies initial conditions.

We remind that $\Theta(0) = I$ and even $\Theta(np) = I$ for all $n = 0, 1, 2, 3, \dots$ because

$$\Phi(np, 0) = [\Phi(p, 0)]^n = \exp((np)F)$$

$\Phi(t, 0)$ is piecewise continuously differentiable or continuously differentiable depending on if $A(t)$ is piecewise continuous or continuous. Therefore $\Theta(t)$ has the same property because $\exp\left(-\frac{t}{p}G\right)$ is continuously differentiable. $\Theta(t)$ is also invertible for all t as a product of two invertible matrices $\Phi(t, 0)$ and $\exp(-tF)$.

We check now that $\Theta(t)$ is p -periodic, namely that $\Theta(t+p) = \Theta(t)$ for all $t \in \mathbb{R}$.

$$\begin{aligned} \Theta(t+p) &\stackrel{def}{=} \Phi(t+p, 0) \exp(-(t+p)F) \\ &= \Phi(t+p, 0) \exp(-pF) \exp(-tF) = \Phi(t+p, 0) \overbrace{\exp(-G)}^{(\exp(G))^{-1}} \exp(-tF) \\ &\quad \underbrace{(\Phi(p,0))^{-1} = \Phi(0,p)} \\ &= \Phi(t+p, 0) \overbrace{(\exp(G))^{-1}}^{\Phi(p,p)=I} \exp(-tF) \end{aligned}$$

We remind that $\exp(G) = \exp(\log(\Phi(p, 0))) = \Phi(p, 0)$, therefore, by Chapman-Kolmogorov relations $\exp(-G) = (\exp(G))^{-1} = \Phi(p, 0)^{-1} = \Phi(0, p)$. Therefore, using the main relation for the monodromy matrix (??) $\Phi(t+p, 0) = \Phi(t, 0)\Phi(p, 0)$ together with the relation $\exp(-G) = \Phi(0, p)$, we arrive to

$$\Theta(t+p) = \Phi(t, 0) \overbrace{\Phi(p, 0)\Phi(0, p)}^{\Phi(p,p)=I} \exp(-tF) = \Phi(t, 0) (I) \exp(-tF) \stackrel{def}{=} \Theta(t),$$

where we also used that $\Phi(p, 0)\Phi(0, p) = I$ in the last step. Therefore $\Theta(t)$ is periodic with period p . ■

1.1 Logarithm of a matrix. Existence and calculation.

We will formulate a theorem and give a proof to it (simpler than in the book) about the existence of a matrix logarithm.

Definition

The matrix G is a **logarithm of matrix** H or $G = \log(H)$ if $\exp(G) = \exp(\log(H)) = H$.

Consider a nonsingular matrix H and it's a canonical Jordan form J :

$$H = TJT^{-1}$$

where T is invertible matrix. Then if there is $Q \in \mathbb{C}^{N \times N}$, such that $\exp(Q) = J$ it means that

$$Q = \log(J), \quad J = \exp(Q)$$

then according to the properties of the exponent of similar matrices, and the definition of the matrix logarithm

$$\begin{aligned} H &= TJT^{-1} = T \exp(Q)T^{-1} = T \exp(\log(J))T^{-1} = \\ &= \exp(T \log(J)T^{-1}) \stackrel{\text{def}}{=} \exp(\log(H)) \end{aligned}$$

and

$$\log(H) = T \log(J)T^{-1}$$

where we used the relation for exponent for similar matrices: that if $A = TBT^{-1}$ then $\exp(A) = T \exp(B)T^{-1}$.

It means that to calculate the logarithm of an arbitrary matrix H it is enough to calculate the logarithm of it's Jordan canonical form. For $H = TJT^{-1}$

$$\log(H) = T \log(J)T^{-1}$$

Definition.

We say that G is a principal logarithm $G = \text{Log}(H)$ of the matrix H if G is a matrix logarithm of H and

$$\begin{aligned} \sigma(H) &= \{\exp(\lambda) : \lambda \in \sigma(G)\} \\ \sigma(G) &= \{\text{Log}(\mu) : \mu \in \sigma(H)\} \end{aligned}$$

where $\text{Log}(\mu)$ is the scalar principal logarithm:

$$z = e^{\text{Log}(z)}; \quad \arg(\text{Log}(z)) = \text{Im}(\text{Log}(z)) \in (-\pi, \pi].$$

This definition implies the explicit one to one correspondence between eigenvalues to H and eigenvalues to G . Essentially the second relation is non-trivial.

Theorem. Proposition 2.29, p. 53.

If $H \in \mathbb{C}^{N \times N}$ is invertible, then there exists a principle logarithm $\text{Log}(H)$.

Proof.

We have established above that it is enough to investigate existence of logarithm for the similar canonical Jordan form J of the matrix H . So without loss of generality we may assume that H is a canonical Jordan form J . Exponent of a Jordan matrix consists of exponents of it's blocks. Therefore it is enough to establish the existence of logarithm for each Jordan block J_j in J , $j = 1, \dots, s$ where s is the number of distinct eigenvalues to H and J_j has size $n_j \times n_j$

$$J_j = \begin{bmatrix} \lambda_j & 1 & 0 & \dots & 0 & 0 \\ 0 & \lambda_j & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & \lambda_j & 1 \\ 0 & 0 & 0 & \dots & 0 & \lambda_j \end{bmatrix}$$

$J_j = \lambda_j \left(I + \frac{1}{\lambda_j} \mathcal{N}_j \right)$ where

$$\mathcal{N}_j = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}$$

From the classical Maclaurin series for $\log(1+x) = \sum_{p=1}^{\infty} \frac{(-1)^{p+1}}{p} x^p$ valid for $|x| < 1$, and for exp we get

$$\exp(\log(1+x)) = 1+x$$

We will try to calculate $\log(J_j) = \log\left(\lambda_j I \left(I + \frac{1}{\lambda_j} \mathcal{N}_j\right)\right) = \log(\lambda_j)I + \log\left(I + \frac{1}{\lambda_j} \mathcal{N}_j\right)$

Point out that we use the condition that $\lambda_j \neq 0!!!$

We formally write the Maclaurin series for $\log\left(I + \frac{1}{\lambda_j} \mathcal{N}_j\right)$:

$$\log\left(I + \frac{1}{\lambda_j} \mathcal{N}_j\right) = \sum_{p=1}^{n_j-1} \frac{(-1)^{p+1}}{p} \left(\frac{1}{\lambda_j} \mathcal{N}_j\right)^p$$

and observe that the Maclaurin series for $\log\left(1 + \frac{1}{\lambda_j} \mathcal{N}_j\right)$ is a **finite sum** because all larger powers of \mathcal{N}_j in the series cancel. We have therefore that

$$\exp\left(\log\left(I + \frac{1}{\lambda_j} \mathcal{N}_j\right)\right) = I + \frac{1}{\lambda_j} \mathcal{N}_j$$

and

$$\begin{aligned} & \exp(\log(\lambda_j)I) \exp\left(\log\left(I + \frac{1}{\lambda_j}\mathcal{N}_j\right)\right) = \\ \exp\left(\log(\lambda_j)I + \log\left(I + \frac{1}{\lambda_j}\mathcal{N}_j\right)\right) &= \lambda_j\left(I + \frac{1}{\lambda_j}\mathcal{N}_j\right) = J_j \end{aligned}$$

We define

$$G_j \stackrel{\text{def}}{=} \log(\lambda_j)I + \sum_{p=1}^{n_j-1} \frac{(-1)^{p+1}}{p} \left(\frac{1}{\lambda_j}\mathcal{N}_j\right)^p$$

Now we check that this expression G_j is actually a matrix logarithm $G_j = \log(J_j)$ for the Jordan block J_j by just checking that it satisfies the definition of the matrix logarithm. Point out that the diagonal matrix $\log(\lambda_j)I$ commutes with any matrix. Therefore applying formula $\exp(\log(1+x)) = 1+x$ for series for $\exp(x)$ and $\log(1+x)$ to similar converging series of commuting matrices we arrive to the desired relation.

$$\begin{aligned} \exp(G_j) &= \exp\left(\log(\lambda_j)I + \sum_{p=1}^{n_j-1} \frac{(-1)^{p+1}}{p} \left(\frac{1}{\lambda_j}\mathcal{N}_j\right)^p\right) \\ &= \exp(\log(\lambda_j)I) \exp\left(\sum_{p=1}^{n_j-1} \frac{(-1)^{p+1}}{p} \left(\frac{1}{\lambda_j}\mathcal{N}_j\right)^p\right) \\ &= \exp(\log(\lambda_j)I) \exp\left(\log\left(I + \frac{1}{\lambda_j}\mathcal{N}_j\right)\right) = (\lambda_j I) \left(I + \frac{1}{\lambda_j}\mathcal{N}_j\right) = J_j \end{aligned}$$

In the Jordan canonical form J eigenvalues stand on diagonal and are easy to control. All calculations that we have carried out are correct because $\lambda_j \neq 0$. We can choose logarithms $\log(\lambda_j)$ in these calculations as principle values of logarithm $\text{Log}(\lambda_j)$. In this case the logarithm of J_j will be principal logarithm, because there will be one to one correspondence between eigenvalues λ_j to J_j and eigenvalues $\text{Log}(\lambda_j)$ to $\text{Log}(J_j)$ that are diagonal elements in corresponding matrices. They will have the same algebraic multiplicity and the same geometric multiplicity 1 (one linearly independent eigenvector for each Jordan block)

Therefore the existence of the principal logarithm is established also for J and for H , that is a matrix similar to J . The same correspondence as above is valid for the eigenvalues to H and to $\text{Log}(H)$ because eigenvalues to similar matrices H and J are the same. The number of linearly independent eigenvectors corresponding to each distinct eigenvalue (geometric multiplicity) will be also the same. ■

1.2 Floquet multipliers and exponents and bounds of solutions to periodic systems. equations.

Definition.

Eigenvalues of the monodromy matrix $\Phi(p, 0)$ are called **Floquet's multipliers** or **characteristic multipliers**.

A Floquet multiplier is called semisimple if it is semisimple as an eigenvalue to the monodromy matrix $\Phi(p, 0)$.

Definition.

Eigenvalues of the logarithm of the monodromy matrix are called **Floquet's exponents** or **characteristic exponents**.

Theorem 2.31, p.54 on boundedness and zero limits of solutions to periodic linear systems.

1) Every solution to a periodic linear system of ODEs is bounded on \mathbb{R}_+ if and only if the absolute value of each Floquet multiplier is not greater than 1 and any Floquet multiplier with absolute value 1 is semisimple.

2) Every solution to a periodic linear system of ODEs tends to zero at $t \rightarrow \infty$ if and only if the absolute value of each Floquet multiplier is strictly less than 1.

Proof. (required at the exam)

By the Floquet theorem any solution $x(t)$ to system

$$x'(t) = A(t)x(t), \quad A(t+p) = A(t), \quad \forall t \in \mathbb{R} \quad (21)$$

satisfying initial conditions

$$x(\tau) = \xi$$

is represented as

$$\begin{aligned} x(t) &= \Phi(t, \tau)\xi = \overbrace{\Theta(t) \exp(tF)}^{\Phi(t,0)} \Phi(0, \tau)\xi = \Theta(t) \overbrace{\exp(tF)\zeta}^{y(t)} \\ &= \Theta(t)y(t) \end{aligned}$$

where

$$F = \frac{1}{p} \text{Log} \left(\begin{matrix} \text{monodromy matrix} \\ \Phi(p, 0) \end{matrix} \right), \quad \zeta = \Phi(0, \tau)\xi.$$

$\Theta(t)$ is a p -periodic continuous or piecewise continuous matrix valued function. $\Theta(t)$ is invertible for all t .

We define $y(t) = \exp(tF)\zeta$ as a solution to the I.V:P. for an autonomous linear equation:

$$y'(t) = F y, \quad y(0) = \zeta \quad (22)$$

$y(t) = \Theta^{-1}(t)x(t)$, and $x(t) = \Theta(t)y(t)$. The mapping $\Theta(t)$ determines a one to one correspondence

between solutions $x(t)$ to the periodic system (21) and solutions $y(t)$ to the autonomous system (22). The periodicity and continuity properties of $\Theta(t)$ and $\Theta^{-1}(t)$ imply that there is a constant $M > 0$ such that $\|\Theta(t)\| \leq M$ and $\|\Theta^{-1}(t)\| \leq M$ for all $t \in \mathbb{R}$. It implies that

$$\|x(t)\| \leq M \|y(t)\| \quad \text{and} \quad \|y(t)\| \leq M \|x(t)\|$$

Therefore

- 1) $\|x(t)\|$ is bounded on \mathbb{R}_+ if and only if corresponding $\|y(t)\| = \|\exp(tF)\zeta\|$ is bounded on \mathbb{R}_+ .
- 2) $\|x(t)\| \rightarrow 0$ when $t \rightarrow \infty$ if and only if corresponding $\|y(t)\| \rightarrow 0$ when $t \rightarrow \infty$.

Since $\text{Log}(\Phi(p, 0)) = G = pF$, and $\Phi(p, 0) = \exp(pF)$ it follows that

$$\begin{aligned} \sigma(\Phi(p, 0)) &= \{\exp(\lambda p) : \lambda \in \sigma(F)\} \\ \sigma(F) &= \left\{ \frac{1}{p} \text{Log}(\mu) : \mu \in \sigma(\Phi(p, 0)) \right\} \end{aligned}$$

and that algebraic and geometric multiplicities of each $\lambda \in \sigma(F)$ coincide with those of $\exp(p\lambda) \in \sigma(\Phi(p, 0))$.

The following connections between the properties of Floquet multipliers and properties of corresponding eigenvalues to the matrix $F = \frac{1}{p} \text{Log}(\Phi(p, 0))$ are direct consequences of the Euler formula for the complex exponent and properties of complex logarithm:

$$\begin{aligned} \text{Log}(z) &= \ln(|z|) + i \text{Arg}(z) \\ \exp(z) &= \exp(\text{Re } z)(\cos(\arg z) + i \sin(\arg z)) \\ |\exp(z)| &= \exp(\text{Re } z) < 1 \iff \text{Re } z < 0 \\ |\exp(z)| &= \exp(\text{Re } z) \leq 1 \iff \text{Re } z \leq 0 \\ |\exp(z)| &= \exp(\text{Re } z) = 1 \iff \text{Re } z = 0 \end{aligned}$$

a) The Floquet multiplier $\mu \in \sigma(\Phi(p, 0))$, has $|\mu| < 1$ if and only if $\text{Re } \text{Log}(\mu) < 0$ that is if the corresponding eigenvalue $\lambda = \frac{1}{p} \text{Log}(\mu)$ to F has $\text{Re } \text{Log}(\mu) < 0$.

b) The Floquet multiplier $\mu \in \sigma(\Phi(p, 0))$, has $|\mu| \leq 1$ if and only if $\text{Re } \text{Log}(\mu) \leq 0$ that is if the corresponding eigenvalue $\lambda = \frac{1}{p} \text{Log}(\mu)$ to F has $\text{Re } \text{Log}(\mu) \leq 0$.

c) The Floquet multiplier $\mu \in \sigma(\Phi(p, 0))$, with $|\mu| = 1$ is semisimple if and only if the corresponding eigenvalue $\lambda = \frac{1}{p} \text{Log}(\mu)$ to F having $\text{Re } \text{Log}(\mu) = 0$ is semisimple.

Known relations between properties of solutions to an autonomous system and the spectrum of corresponding matrix applied to the system $y'(t) = Fy$ and to the spectrum $\sigma(F)$ of the matrix F together with statements 1), 2), a), b), c) in the present proof imply the statement of the theorem. ■

Proposition 2.20. p. 45. On periodic solutions to periodic linear systems

The system $x'(t) = A(t)x(t)$ with p - periodic $A(t) = A(t + p)$ has a non-zero p - periodic solution if and only if the monodromy matrix $\Phi(p, 0)$ has an eigenvalue $\lambda = 1$. A more general statement is also valid.

The system has a non-zero np - periodic solution for $n \in \mathbb{N}$ if and only if the monodromy matrix $\Phi(p, 0)$ has an eigenvalue λ such that $\lambda^n = 1$. \square

Proof. Consider an eigenvector v corresponding to the eigenvalue λ . Then $v \neq 0$, $\Phi(p, 0)v = \lambda v$ and

$$[\Phi(p, 0)]^n v = \lambda^n v = 1 \cdot v = v$$

We will show that the solution to the system, with initial data $x(0) = v$ has period np . This solution is given by the transition matrix: $x(t) = \Phi(t, 0)v$. Using this representation and applying the factorisation property of transition matrices for periodic systems we arrive to

$$x(t + np) = \Phi(t + np, 0)v = \Phi(t, 0) [\Phi(p, 0)]^n v = \Phi(t, 0)v = x(t), \quad \forall t \in \mathbb{R}$$

It shows that $x(t)$ is periodic with period np .

Supposing that there is a periodic solution $x(t + np) = x(t)$ and repeating the same calculation backwards we arrive to that $x(0) = v$ is an eigenvector corresponding to an eigenvalue λ such that $\lambda^n = 1$.

Carry out this backward argument as an exercise!

Lecture 13

Examples and exercises

on periodic and general linear systems of ODEs.

1. Monodromy matrix: $\Phi(p, 0)$, where p is a period of coefficients in the matrix $A(t) = A(t + p)$.
2. If all eigenvalues λ to $\Phi(p, 0)$ have $|\lambda| < 1$, then all solutions to $x'(t) = A(t)x(t)$ tend to zero with $t \rightarrow \infty$.
3. If all eigenvalues λ to $\Phi(p, 0)$ have $|\lambda| \leq 1$, and eigenvalues with $|\lambda| = 1$ are semisimple, then all solutions to $x'(t) = A(t)x(t)$ are bounded.

Comment conditions in 2. ,3. are also necessary !

Corollary 2.33, p. 59

We consider a periodic linear system $x'(t) = A(t)x(t)$, $A(t+p) = A(t)$.

If $\int_0^p \text{tr}(A(s)ds)$ has a positive real part, then the equation has at least one solution $x(t)$ that is unbounded, or expressing it more explicitly, the upper limit of it's norm is infinite: $\limsup_{t \rightarrow \infty} \|x(t)\| = \infty$.

□

Proof.

We remind that the transition matrix $\Phi(t, \tau)$ satisfies the initial value problem:

$$\begin{aligned} \frac{d}{dt}\Phi(t, \tau) &= A(t)\Phi(t, \tau) \\ \Phi(\tau, \tau) &= I \end{aligned}$$

Arbitrary solution to the initial problem $x'(t) = A(t)x(t)$, $x(\tau) = \xi$ will be expressed as

$$x(t) = \Phi(t, \tau)\xi$$

According to Abel - Liouville's formula and considerations before

$$\det(w(t, 0)) = \det(w(0, 0)) \exp\left(\int_0^t \text{tr}(A(s)ds)\right)$$

It implies for $\Phi(t, 0)$

$$\begin{aligned} |\det(\Phi(t, 0))| &= \left| \det(\Phi(0, 0)) \exp\left(\int_0^t \text{tr}(A(s)ds)\right) \right| = \\ \left| \exp\left(\int_0^t \text{tr}(A(s)ds)\right) \right| &= \left| \exp\left(\text{Re}\left(\int_0^t \text{tr}(A(s)ds)\right)\right) \right| \end{aligned}$$

Therefore, for $t = p$ we get that if $\text{Re}\left(\int_0^p \text{tr}(A(s)ds)\right) > 0$ then

$$|\det(\Phi(p, 0))| = \left| \exp\left(\text{Re}\int_0^p \text{tr}(A(s)ds)\right) \right| > 1.$$

On the other hand $\det(\Phi(p, 0))$ is a product of eigenvalues μ_k to the monodromy matrix $\Phi(p, 0)$ with multiplicities m_k

It follows from the structure of the similar Jordan matrix

$$\begin{aligned} \Phi(p, 0) &= T^{-1}JT \\ \det \Phi(p, 0) &= \det(T)^{-1} \det(T) \det(J) = \det(J) \end{aligned}$$

$$|\det(\Phi(p, 0))| = \prod_{k=1}^s |\mu_k|^{m_k} > 1$$

To have this product greater than 1 we must have at least one eigenvalue μ_p with $|\mu_p| > 1$. According to one of Floquet theorems, all solutions to a periodic system of ODEs are bounded if and only if all eigenvalues to the monodromy matrix have absolute value $|\lambda| \leq 1$ and those with $|\lambda| = 1$ are semisimple. Therefore existence of an eigenvalue with $|\lambda| > 1$ implies existence of a solution $x(t)$ that is not bounded and therefore $\limsup_{t \rightarrow \infty} \|x(t)\| = \infty$. ■

For example we can choose the initial condition $x(0) = v_p$ with v_p being the eigenvector to $\Phi(p, 0)$ corresponding to the eigenvalue $|\mu_p| > 1$. Then the solution

$$\begin{aligned} x(t) &= \Phi(t, 0)v_p \\ \text{Let } t_n &= np \\ \Phi(np, 0)v_p &= [\Phi(p, 0)]^n v_p = (\mu_p)^n v_p \end{aligned}$$

with $|\mu_p| > 1$. Therefore $x(t_n) = (\mu_p)^n v_p$, and

$$\|x(t_n)\| = |\mu_p|^n \|v_p\| \xrightarrow{n \rightarrow \infty} \infty$$

and $\limsup_{t \rightarrow \infty} \|x(t)\| = \infty$.

□

We give also a geometric interpretation of this result. Consider a unite cube build on standard base vectors e_1, \dots, e_N at time $t = 0$. Consider how the volume $\text{Vol}(t)$ of this cube changes under the action of the linear transformation by the transition matrix $\Phi(t, 0)$ of our periodic system. Point out that $I = [e_1, \dots, e_N]$. It implies that the figure of interest is the parallelepiped build on columns of the transition matrix $\Phi(t, 0)$. One of the main properties of periodic system is that $\Phi(np, 0) = [\Phi(p, 0)]^n$. Therefore

$$\text{Vol}(np) = |\det([\Phi(p, 0)]^n)| = |\det([\Phi(p, 0)])|^n = \left[\exp \left(\text{Re} \left(\int_0^p \text{tr}(A(s)ds) \right) \right) \right]^n$$

If $\text{Re} \left(\int_0^p \text{tr}(A(s)ds) \right) > 0$ then $\exp \left(\text{Re} \left(\int_0^p \text{tr}(A(s)ds) \right) \right) > 1$. It implies that

$$\lim_{n \rightarrow \infty} \text{Vol}(np) = \infty$$

Therefore along the sequence of times $\{t = np, \quad n = 1, 2, 3, \dots\}$ we observe that $\text{Vol}(np)$ is unbounded. It implies also that

$$\limsup_{t \rightarrow \infty} \|\text{Vol}(t)\| = \infty$$

The fact that $\lim_{n \rightarrow \infty} \text{Vol}(np) = \infty$ implies that the diameter $D(np)$ of the parallelepiped build on columns of $\Phi(np, 0)$ calculated at these discrete time points, is also unbounded $\lim_{n \rightarrow \infty} D(np) = \infty$. It in turn means that there should be a solution that has the property $\lim_{n \rightarrow \infty} \|x(np)\| = \infty$. □