

TMA683 Tillämpad matematik

Övningsuppgifter (boken FEM)

28 oktober 2021

This document contains the exercises from the compendium from M. Asadzadeh (23.08.2018). Particularly relevant exercises are marked with (*).

Propositions or hints for solutions are given at the end of the file (thanks to Sebastian Persson).

Thank you for reporting typos or errors via [email](#).

1. CHAPTER 4: POLYNOMIAL APPROXIMATION IN $1d$

4.1 Prove that $V_0^{(q)} = \{v \in \mathcal{P}^{(q)}(0, 1), v(0) = 0\}$ is a subspace of $\mathcal{P}^{(q)}(0, 1)$.

4.3 Consider the ODE

$$\dot{u}(t) = u(t), \quad 0 < t < 1, \quad u(0) = 1.$$

Compute its Galerkin approximation in $\mathcal{P}^{(q)}(0, 1)$ for $q = 1, 2, 3, 4$.

4.4 (*) Compute the stiffness matrix and load vector in a finite element approximation of the BVP

$$-u''(x) = f(x), \quad 0 < x < 1, \quad u(0) = u(1) = 0$$

with $f(x) = x$ and $h = 1/4$.

4.5 We want to find a solution approximation $U(x)$ to

$$-u''(x) = 1, \quad 0 < x < 1, \quad u(0) = u(1) = 0,$$

using the ansatz $U(x) = A \sin(\pi x) + B \sin(2\pi x)$.

- (a) Calculate the exact solution $u(x)$.
- (b) Write down the residual $R(x) = -U''(x) - 1$.
- (c) Use the orthogonality condition

$$\int_0^1 R(x) \sin(n\pi x) dx = 0, \quad n = 1, 2$$

to determine the constants A and B .

- (d) Plot the error $e(x) = |u(x) - U(x)|$.

4.6 Consider the BVP

$$-u''(x) + u(x) = x, \quad 0 < x < 1, \quad u(0) = u(1) = 0.$$

- (a) Verify that the exact solution to the above problem reads

$$u(x) = x - \frac{\sinh(x)}{\sinh(1)}.$$

- (b) Let $U(x)$ be a solution approximation defined by

$$U(x) = A \sin(\pi x) + B \sin(2\pi x) + C \sin(3\pi x),$$

where A, B, C are unknown constants. Compute the residual

$$R(x) = -U''(x) + U(x) - x.$$

- (c) Use the orthogonality conditions

$$\int_0^1 R(x) \sin(n\pi x) dx = 0, n = 1, 2, 3$$

to determine the constants A, B, C .

- 4.7 Let $U(x) = \zeta_0 \phi_0(x) + \zeta_1 \phi_1(x)$ be a solution approximation to

$$-u''(x) = x - 1, \quad 0 < x < \pi, \quad u'(0) = u(\pi) = 0,$$

where ζ_0 and ζ_1 are unknown coefficients and $\phi_0(x) = \cos(\frac{x}{2})$, $\phi_1(x) = \cos(\frac{3x}{2})$.

- (a) Find the analytical solution $u(x)$.
(b) Define the residual $R(x)$.
(c) Compute the constants ζ_0 and ζ_1 using the orthogonality conditions

$$\int_0^\pi R(x) \phi_i(x) dx = 0, i = 0, 1.$$

I.e. by projecting $R(x)$ onto the vector space spanned by $\phi_0(x)$ and $\phi_1(x)$.

- 4.8 Use the projection technique of the previous exercise to solve

$$-u''(x) = 0, \quad 0 < x < \pi, \quad u(0) = 0, u(\pi) = 2,$$

with $U(x) = A \sin(x) + B \sin(2x) + C \sin(3x) + \frac{2}{\pi^2} x^2$ and using the test functions $\{\sin(x), \sin(2x), \sin(3x)\}$.

2. CHAPTER 5: INTERPOLATION, NUMERICAL INTEGRATION IN 1d

5.1 Consider two real numbers $a < b$. By definition of Lagrange's polynomials, one has

$$\lambda_a(x) = \frac{b-x}{b-a} \quad \text{and} \quad \lambda_b(x) = \frac{x-a}{b-a}.$$

Show that

$$\lambda_a(x) + \lambda_b(x) = 1 \quad \text{and} \quad a\lambda_a(x) + b\lambda_b(x) = x.$$

Give a geometric interpretation by plotting $\lambda_a(x)$, $\lambda_b(x)$, $\lambda_a(x) + \lambda_b(x)$ and $a\lambda_a(x)$, $b\lambda_b(x)$, $a\lambda_a(x) + b\lambda_b(x)$.

5.2 (*) Consider the following functions defined for $x \in [0, 1]$:

$$f(x) = x^2 \quad \text{and} \quad g(x) = \sin(\pi x).$$

Find their linear interpolants, denoted by $\Pi f \in \mathcal{P}(0, 1)$, resp. $\Pi g \in \mathcal{P}(0, 1)$. In the same figure, plot f and Πf , as well as g and Πg .

5.3 Determine the linear interpolant of the function, defined for $x \in [-\pi, \pi]$,

$$f(x) = \frac{1}{\pi^2}(x - \pi)^2 - \cos^2(x - \frac{\pi}{2}),$$

where the interval $[-\pi, \pi]$ is divided into 4 equal subintervals.

5.15 Prove that

$$\int_{x_0}^{x_1} f'(\frac{x_0 + x_1}{2})(x - \frac{x_0 + x_1}{2}) dx = 0.$$

5.16 (*) Prove that

$$\begin{aligned} \left| \int_{x_0}^{x_1} f(x) dx - f(\frac{x_0 + x_1}{2})(x_1 - x_0) \right| &\leq \frac{1}{2} \max_{[x_0, x_1]} |f''(x)| \int_{x_0}^{x_1} (x - \frac{x_0 + x_1}{2})^2 dx \\ &\leq \frac{1}{24} (x_1 - x_0)^3 \max_{[x_0, x_1]} |f''(x)|. \end{aligned}$$

Hint: Use a Taylor expansion of f about $x = \frac{x_0 + x_1}{2}$.

3. CHAPTER 7: TWO-POINT BOUNDARY VALUE PROBLEMS

7.1 Consider the two-point BVP

$$-u''(x) = f(x), \quad 0 < x < 1, \quad u(0) = u(1) = 0.$$

Let $V = \{v: \|v\| + \|v'\| < \infty, v(0) = v(1) = 0\}$ where $\|\cdot\|$ denotes the L_2 -norm.

- (a) Use V to derive a variational formulation for the above BVP.
- (b) Discuss why V is valid as a vector space of test functions.
- (c) Classify which of the following functions are admissible test functions:

$$\sin(\pi x), \quad x^2, \quad x \ln(x), \quad e^x - 1, \quad x(1 - x).$$

7.3 Consider the two-point BVP

$$-u''(x) = 1, \quad 0 < x < 1, \quad u(0) = u(1) = 0.$$

Let $\mathcal{T}_h : x_j = \frac{j}{4}, j = 0, 1, 2, 3, 4$ denote a partition of the interval $0 < x < 1$ into four subintervals of equal length $h = 1/4$. Let V_h be the corresponding space of continuous piecewise linear functions vanishing at $x = 0$ and $x = 1$.

- (a) Compute a finite element approximation $U \in V_h$ to the above BVP.
- (b) Prove that $U \in V_h$ is unique.

7.5 (*) Consider the two-point BVP, for $x \in I = (0, 1)$:

$$\begin{aligned} -(a(x)u'(x))' &= f(x) \\ u(0) &= 0, \quad a(1)u'(1) = g_1, \end{aligned}$$

where a is a positive function and g_1 a constant.

- (a) Derive the variational formulation of the above problem.
- (b) Discuss how the boundary conditions are implemented.

7.6 Consider the two-point BVP, for $x \in I = (0, 1)$,

$$\begin{aligned} -u''(x) &= 0 \\ u(0) &= 0, \quad u'(1) = 7. \end{aligned}$$

Divide the interval I into two subintervals of length $h = \frac{1}{2}$. Let V_h be the corresponding space of continuous piecewise linear functions vanishing at $x = 0$.

- (a) Formulate a finite element method for the above problem.
- (b) Calculate by hand the finite element approximation $U \in V_h$ to the above BVP.
- (c) Study how the boundary condition at $x = 1$ is approximated.

7.7 (*) Consider the two-point BVP

$$-u''(x) = 0, \quad 0 < x < 1, \quad u'(0) = 5, u(1) = 0.$$

Let $\mathcal{T}_h : x_j = \frac{j}{N}, j = 0, 1, \dots, N, h = 1/N$ denote a uniform partition of the interval $0 < x < 1$ into N subintervals. Let V_h be the corresponding space of continuous piecewise linear functions.

- (a) Use V_h , with $N = 3$, and formulate a finite element method for the above problem.
- (b) Compute the finite element approximation $U \in V_h$ assuming $N = 3$.

7.8 Consider the problem of finding a solution approximation to

$$-u''(x) = 1, \quad 0 < x < 1, \quad u'(0) = u'(1) = 0.$$

Let \mathcal{T}_h be a partition of the interval $0 < x < 1$ into two subintervals of equal length $h = \frac{1}{2}$. Let V_h be the corresponding space of continuous piecewise linear functions.

- (a) Can you find an exact solution to the above problem by integrating twice?
- (b) Compute a finite element approximation $U \in V_h$ to u if possible.

7.11 Consider the finite element method applied to

$$-u''(x) = 0, \quad 0 < x < 1, \quad u(0) = \alpha, u'(1) = \beta,$$

where α and β are given constants. Assume that the interval $[0, 1]$ is divided into three subintervals of equal length $h = 1/3$ and that $\{\varphi_j\}_{j=0}^3$ is a nodal basis of V_h , the corresponding space of continuous piecewise linear functions.

- (a) Verify that the ansatz

$$U(x) = \alpha\varphi_0(x) + \zeta_1\varphi_1(x) + \zeta_2\varphi_2(x) + \zeta_3\varphi_3(x),$$

yields the following system of equations

$$(1) \quad \frac{1}{h} \begin{pmatrix} -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} \alpha \\ \zeta_1 \\ \zeta_2 \\ \zeta_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \beta \end{pmatrix}.$$

- (b) If $\alpha = 2$ and $\beta = 3$ show that (1) can be reduced to

$$\frac{1}{h} \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} \zeta_1 \\ \zeta_2 \\ \zeta_3 \end{pmatrix} = \begin{pmatrix} 2h^{-1} \\ 0 \\ 3 \end{pmatrix}.$$

- (c) Solve the above system of equation to find $U(x)$.

7.13 Consider the following boundary value problem

$$-au''(x) + bu(x) = 0, \quad 0 \leq x \leq 1, \quad u(0) = u'(1) = 0,$$

where $a, b > 0$ are constants. Let $\mathcal{T}_h : 0 = x_0 < x_1 < \dots < x_N = 1$, be a non-uniform partition of the interval $0 \leq x \leq 1$ into N intervals of length $h_i = x_i - x_{i-1}$, $i = 1, 2, \dots, N$. Let V_h be the corresponding space of continuous piecewise linear functions. Compute the stiffness and mass matrices.

7.14 Show that the FEM with mesh size h for the problem

$$\begin{cases} -u''(x) = 1 & 0 < x < 1 \\ u(0) = 7, u'(1) = 0, \end{cases}$$

with $U(x) = 7\varphi_0(x) + U_1\varphi_1(x) + \dots + U_m\varphi_m(x)$ leads to the linear system of equations $\tilde{A}\tilde{U} = \tilde{b}$, where $\tilde{A} \in \mathbb{R}^{m \times (m+1)}$, $\tilde{U} \in \mathbb{R}^{(m+1) \times 1}$, $\tilde{b} \in \mathbb{R}^{m \times 1}$ are given by

$$\tilde{A} = \frac{1}{h} \begin{pmatrix} -1 & 2 & -1 & 0 & \dots & 0 \\ 0 & -1 & 2 & -1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 & \dots & 0 \end{pmatrix}, \tilde{U} = \begin{pmatrix} 7 \\ U_1 \\ \vdots \\ U_m \end{pmatrix}, \tilde{b} = \begin{pmatrix} h \\ \vdots \\ h \\ h/2 \end{pmatrix}.$$

The above reduces to $AU = b$, with

$$A = \frac{1}{h} \begin{pmatrix} 2 & -1 & 0 & \dots & 0 \\ -1 & 2 & -1 & \dots & 0 \\ \dots & \dots & -1 & 2 & -1 \\ 0 & 0 & \dots & -1 & 2 \end{pmatrix}, U = \begin{pmatrix} U_1 \\ \vdots \\ U_m \end{pmatrix}, b = \begin{pmatrix} h + \frac{7}{h} \\ \vdots \\ h \\ h/2 \end{pmatrix}.$$

4. CHAPTER 8: SCALAR INITIAL VALUE PROBLEMS

8.5a) Compute the solution of

$$\dot{u}(t) + a(t)u(t) = t^2, \quad 0 < t < T, \quad u(0) = 1,$$

where $a(t) = 4$.

5. CHAPTER 9: INITIAL BOUNDARY VALUE PROBLEMS IN 1d

9.7 Consider the inhomogeneous problem

$$\begin{cases} u_t(x, t) - \varepsilon u_{xx}(x, t) = f(x, t), & 0 < x < 1, t > 0 \\ u(0, t) = u_x(1, t) = 0, & t > 0 \\ u(x, 0) = u_0(x), & 0 < x < 1. \end{cases}$$

Show that for the corresponding stationary problem, $u_t = 0$, one has

$$\|u_x\| \leq \frac{1}{\varepsilon} \|f\|.$$

9.13 Consider the wave equation

$$\begin{cases} u_{tt}(x, t) - u_{xx}(x, t) = 0, & x \in \mathbb{R}, t > 0 \\ u(x, 0) = u_0(x), & x \in \mathbb{R} \\ u_t(x, 0) = v_0(x), & x \in \mathbb{R}. \end{cases}$$

Plot the graph of $u(x, 2)$ in the following cases:

(a) $v_0 = 0$ and

$$u_0(x) = \begin{cases} 1, & x < 0 \\ 0, & x > 0. \end{cases}$$

(b) $u_0 = 0$ and

$$v_0(x) = \begin{cases} -1, & -1 < x < 0 \\ 1, & 0 < x < 1 \\ 0, & |x| > 1. \end{cases}$$

6. CHAPTER 4: PROPOSITIONS FOR SOLUTIONS

4.1 Use the definitions of $\mathcal{P}^{(q)}(0, 1)$ and of a subspace.

4.3 Every element $v \in \mathcal{P}^{(q)}(0, 1)$ can be written as

$$v(t) = \sum_{j=0}^q \chi_j t^j.$$

Use this in a VF of the problem.

4.4 See the lecture.

4.5 (a) The exact solution reads $u(x) = \frac{x}{2}(1 - x)$.

(b) The residual reads $R(x) = \pi^2(A \sin(\pi x) + 4B \sin(2\pi x)) - 1$.

(c) $A = \frac{4}{\pi^3}$ and $B = 0$.

4.6 (a) ok

(b)

$$R(x) = (\pi^2 + 1)A \sin(\pi x) + (4\pi^2 + 1)B \sin(2\pi x) + (9\pi^2 + 1)C \sin(3\pi x) - x.$$

(c)

$$A = \frac{2}{\pi(\pi^2 + 1)}, B = -\frac{1}{\pi(4\pi^2 + 1)}, C = \frac{2}{3\pi(9\pi^2 + 1)}$$

4.7 (a)

$$u(x) = \frac{1}{6}(\pi^3 - x^3) + \frac{1}{2}(x^2 - \pi^2)$$

(b)

$$R(x) = -U''(x) - x + 1 = \frac{1}{4}\zeta_0 \cos\left(\frac{x}{2}\right) + \frac{9}{4}\zeta_1 \cos\left(\frac{3x}{2}\right) - x + 1$$

(c)

$$\zeta_0 = 8(2\pi - 6)/\pi, \zeta_1 = \frac{8}{9}\left(\frac{2}{9} - \frac{2}{3}\pi\right)/\pi$$

4.8

$$U(x) = (16 \sin(x) + \frac{16}{27} \sin(3x))/\pi^3 + \frac{2}{\pi^2} x^2$$

7. CHAPTER 5: PROPOSITIONS FOR SOLUTIONS

5.1 Insert the definition of

$$\lambda_a(x) = \frac{b-x}{b-a} \quad \text{and} \quad \lambda_b(x) = \frac{x-a}{b-a}.$$

into

$$\lambda_a(x) + \lambda_b(x) \quad \text{and} \quad a\lambda_a(x) + b\lambda_b(x)$$

to answer the exercise.

5.2 Use the definition of the linear interpolant, see lecture.

5.3

$$\Pi_1 f(x) = \begin{cases} 4 - 11(x + \pi)/(2\pi), & -\pi \leq x \leq -\frac{\pi}{2} \\ 5/4 - (x + \frac{\pi}{2})/(2\pi), & -\frac{\pi}{2} \leq x \leq 0 \\ 1 - 7x/(2\pi), & 0 \leq x \leq \frac{\pi}{2} \\ 3(x - \pi)/(2\pi), & \frac{\pi}{2} \leq x \leq \pi \end{cases}$$

5.15 Observe that the term $f'(\frac{x_0+x_1}{2})$ does not depend on x and use the formula $(a+b)(a-b) = a^2 - b^2$.

5.16 This is the local error of the midpoint rule. Use a Taylor expansion (with rest term) of f about $x = \frac{x_0+x_1}{2}$ to show the exercise.

8. CHAPTER 7: PROPOSITIONS FOR SOLUTIONS

- 7.1 (a) See lecture.
(b) See lecture.
(c) The following functions are admissible test functions:

$$\sin(\pi x), \quad x(1-x).$$

- 7.3 (a) See lecture.
(b) Assume that one has more than one solution to the FE and, using the FE formulation, find a contradiction.
- 7.5 (a) Similar to the lecture.
(b) Consider possible additional terms in the last vector.
- 7.6 (a) Similar to the lecture.
(b) Long computation
(c) tba
- 7.7 (a) Find $u_h \in V_h$ such that

$$\int_0^1 u_h(x)v_h(x) \, dx = -5v_h(0)$$

for all $v \in V_h^0$.

- (b) The FE solution reads

$$u_h(x) = \alpha_0\varphi_0(x) + \alpha_1\varphi_1(x) + \alpha_2\varphi_2(x),$$

where φ_j are the hat functions and $\alpha_0 = -5, \alpha_1 \approx -3.333, \alpha_2 \approx -1.667$.

- 7.8 (a) Integrate the problem twice and do not forget the two integration constants.
(b) Observe that the resulting matrix from a FE discretisation is not invertible.
- 7.11 (a) Observe that one has non-homogeneous Dirichlet BC and hence need two spaces (trial, resp. test)

$$V = \{v: v, v' \in L^2(0,1), v(0) = \alpha\} \quad \text{and} \quad V^0 = \{v: v, v' \in L^2(0,1), v(0) = 0\}$$

for the VF (similarly for the FE formulation).

- (b) ok
(c) One can use matlab to compute such solution.
- 7.13 Similar to the lecture.
- 7.14 Similar to the lecture.

9. CHAPTER 8: PROPOSITIONS FOR SOLUTIONS

8.5a)

$$u(t) = e^{-4t} + \frac{1}{32}(8t^2 - 4t + 1)$$

10. CHAPTER 9: PROPOSITIONS FOR SOLUTIONS

9.7 Recall the definition of the L^2 -norm:

$$\|u\|^2 = (u, u) = \int_0^L u(x)u(x) \, dx$$

and multiply the problem with an appropriate function and integrate (in space). Poincaré inequality could also be of some use.

9.13 One may use d'Alembert's formula ([wiki](#))

$$u(x, t) = \frac{1}{2}(u_0(x - t) - u_0(x + t)) + \frac{1}{2} \int_{x-t}^{x+t} v_0(y) \, dy.$$