

Chapter 7: The heat equation in 1d (summary)

February 5, 2022

Goal: Briefly study the exact solution to some heat equations and present a numerical discretisation.

- Let us start with some **stability estimates** for the following heat equation

$$\begin{cases} u_t(x, t) - u_{xx}(x, t) = f(x, t) & 0 < x < 1, 0 < t \leq T \\ u(0, t) = 0, u_x(1, t) = 0 & 0 < t \leq T \\ u(x, 0) = u_0(x) & 0 < x < 1, \end{cases}$$

where u_0 and f are given functions.

The solution to the above problem satisfy the following estimates

$$\begin{aligned} \|u(\cdot, t)\|_{L^2(0,1)} &\leq \|u_0\|_{L^2(0,1)} + \int_0^t \|f(\cdot, s)\|_{L^2(0,1)} \, ds. \\ \|u_x(\cdot, t)\|_{L^2(0,1)} &\leq \|u'_0\|_{L^2(0,1)} + \int_0^t \|f(\cdot, s)\|_{L^2(0,1)} \, ds. \end{aligned}$$

When $f = 0$, one gets

$$\|u(\cdot, t)\|_{L^2(0,1)} \leq \|u_0\|_{L^2(0,1)} e^{-2t}.$$

When $f = 0$ and some fixed $\varepsilon > 0$, one gets for all $t \in (0, T]$

$$\int_\varepsilon^t \|u_t(\cdot, s)\|_{L^2(0,1)} \, ds \leq \frac{1}{2} \sqrt{\ln\left(\frac{t}{\varepsilon}\right)} \|u_0\|_{L^2(0,1)}.$$

- Next, we **discretise the inhomogeneous heat equation** with homogeneous Dirichlet boundary conditions

$$\begin{cases} u_t(x, t) - u_{xx}(x, t) = f(x, t) & 0 < x < 1, 0 < t \leq T \\ u(0, t) = u(1, t) = 0 & 0 < t \leq T \\ u(x, 0) = u_0(x) & 0 < x < 1, \end{cases}$$

where u_0 and f are given functions.

Since it is seldom possible to find the exact solution $u(x, t)$ to the above problem, we need to find a numerical approximation of it. We proceed as follows

- To get a VF of the heat equation, consider the test/trial space

$V^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v, v' \in L^2(0, 1), v(0) = v(1) = 0\}$. Then, multiply the DE by a test function $v \in V^0$, integrate over $[0, 1]$, and use integration by parts to get the VF: For each $0 < t \leq T$

$$\text{Find } u(\cdot, t) \in V^0 \text{ s.t. } (u_t(\cdot, t), v)_{L^2} + (u_x(\cdot, t), v_x)_{L^2} = (f(\cdot, t), v)_{L^2} \quad \forall v \in V^0 \quad (\text{VF})$$

with the initial condition $u(x, 0) = u_0(x)$.

- To get a FE problem, we consider the following subspace of the above space V^0

$V_h^0 = \{v: [0, 1] \rightarrow \mathbb{R} : v \text{ cont. pw. linear on unif. partition } T_h, v(0) = v(1) = 0\} = \text{span}(\varphi_1, \dots, \varphi_m)$, where $h = \frac{1}{m+1}$ and φ_j are the hat functions.

The FE problem then reads: For each $0 < t \leq T$

$$\text{Find } u_h(\cdot, t) \in V_h^0 \text{ s.t. } (u_{h,t}(\cdot, t), \chi)_{L^2} + (u_{h,x}(\cdot, t), v_{h,x})_{L^2} = (f(\cdot, t), v_h)_{L^2} \quad \forall v_h \in V_h^0 \quad (\text{FE})$$

with the initial condition $u_h(x, 0) = \pi_h u_0(x)$ the cont. pw. linear interpolant of u_0 .

3. From the above FE problem, we obtain a system of linear ODE by choosing the test functions $v_h = \varphi_i$ for $i = 1, \dots, m$ and writing $u_h(x, t) = \sum_{j=1}^m \zeta_j(t) \varphi_j(x)$ with unknown coordinates $\zeta_j(t)$.

Inserting everything in (FE), one gets the ODE

$$\begin{aligned} M\dot{\zeta}(t) + S\zeta(t) &= F(t) \\ \zeta(0), \end{aligned} \tag{ODE}$$

where M is the (already seen) $m \times m$ mass matrix, S is the (already seen) $m \times m$ stiffness matrix, $F(t)$ is an $m \times 1$ vector with entries $F_i(t) = (f(\cdot, t), \varphi_i)_{L^2}$ for $i = 1, \dots, m$, the initial condition is given by

$$\zeta(0) = \begin{pmatrix} u_0(x_1) \\ \vdots \\ u_0(x_m) \end{pmatrix},$$

and the unknown vector reads

$$\zeta(t) = \begin{pmatrix} \zeta_1(t) \\ \vdots \\ \zeta_m(t) \end{pmatrix}.$$

4. To find a numerical approximation of $\zeta(t)$ at some discrete time grid $t_0 = 0 < t_1 < \dots < t_N = T$, with $t_j - t_{j-1} = k = \frac{T}{N}$, one can for instance use backward Euler scheme which reads

$$\begin{aligned} \zeta^{(0)} &= \zeta(0) \\ (M + kS)\zeta^{(n+1)} &= M\zeta^{(n)} + kF(t_{n+1}) \quad \text{for } n = 0, 1, 2, \dots, N-1. \end{aligned}$$

Solving these linear systems at each time step provides numerical approximations $\zeta^{(n)} \approx \zeta(t_n)$ that can be inserted in the FE solution to get approximations to the exact solution to the heat equation $\sum_{j=1}^m \zeta_j(t_n) \varphi_j(x) \approx u(x, t_n)$.

Further resources:

- [heat eq. at wikipedia.org](https://en.wikipedia.org/wiki/Heat_equation)
- [heat eq. at math.lamar.edu](https://math.lamar.edu/)