

# Lecture 9: Feature selection and regularised regression

---

Rebecka Jörnsten, Mathematical Sciences

**MSA220/MVE441** Statistical Learning for Big Data

25<sup>th</sup> April 2022



**CHALMERS**  
UNIVERSITY OF TECHNOLOGY



UNIVERSITY OF GOTHENBURG



# Goals of modelling

---

1. **Predictive strength:** How well can we reconstruct the observed data? Has been most important so far.



# Goals of modelling

---

1. **Predictive strength:** How well can we reconstruct the observed data? Has been most important so far.
2. **Model/variable selection:** Which variables are **part of the true model**? This is about uncovering structure to allow for mechanistic understanding.



# Feature Selection

---



## Remember ordinary least-squares (OLS)

---

Consider the model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$$

where

- ▶  $\mathbf{y} \in \mathbb{R}^n$  is the **outcome**,  $\mathbf{X} \in \mathbb{R}^{n \times (p+1)}$  is the **design matrix**,  $\boldsymbol{\beta} \in \mathbb{R}^{p+1}$  are the **regression coefficients**, and  $\boldsymbol{\varepsilon} \in \mathbb{R}^n$  is the **additive error**



# Remember ordinary least-squares (OLS)

Consider the model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$$

where

- ▶  $\mathbf{y} \in \mathbb{R}^n$  is the **outcome**,  $\mathbf{X} \in \mathbb{R}^{n \times (p+1)}$  is the **design matrix**,  $\boldsymbol{\beta} \in \mathbb{R}^{p+1}$  are the **regression coefficients**, and  $\boldsymbol{\varepsilon} \in \mathbb{R}^n$  is the **additive error**
- ▶ **Five basic assumptions** have to be checked
  - Underlying relationship is linear (1)
  - Zero mean (2), uncorrelated (3) errors with constant variance (4) which are (roughly) normally distributed (5)



## Remember ordinary least-squares (OLS)

Consider the model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$$

where

- ▶  $\mathbf{y} \in \mathbb{R}^n$  is the **outcome**,  $\mathbf{X} \in \mathbb{R}^{n \times (p+1)}$  is the **design matrix**,  $\boldsymbol{\beta} \in \mathbb{R}^{p+1}$  are the **regression coefficients**, and  $\boldsymbol{\varepsilon} \in \mathbb{R}^n$  is the **additive error**
- ▶ **Five basic assumptions** have to be checked
  - Underlying relationship is linear (1)
  - Zero mean (2), uncorrelated (3) errors with constant variance (4) which are (roughly) normally distributed (5)
- ▶ **Centring** ( $\frac{1}{n} \sum_{l=1}^n x_{lj} = 0$ ) and **standardisation** ( $\frac{1}{n} \sum_{l=1}^n x_{lj}^2 = 1$ ) of predictors simplifies interpretation



## Remember ordinary least-squares (OLS)

Consider the model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}$$

where

- ▶  $\mathbf{y} \in \mathbb{R}^n$  is the **outcome**,  $\mathbf{X} \in \mathbb{R}^{n \times (p+1)}$  is the **design matrix**,  $\boldsymbol{\beta} \in \mathbb{R}^{p+1}$  are the **regression coefficients**, and  $\boldsymbol{\varepsilon} \in \mathbb{R}^n$  is the **additive error**
- ▶ **Five basic assumptions** have to be checked
  - Underlying relationship is linear (1)
  - Zero mean (2), uncorrelated (3) errors with constant variance (4) which are (roughly) normally distributed (5)
- ▶ **Centring** ( $\frac{1}{n} \sum_{l=1}^n x_{lj} = 0$ ) and **standardisation** ( $\frac{1}{n} \sum_{l=1}^n x_{lj}^2 = 1$ ) of predictors simplifies interpretation
- ▶ **Centring** the outcome ( $\frac{1}{n} \sum_{l=1}^n y_l = 0$ ) and features removes the need to estimate the intercept



## Feature selection as motivation

---

Analytical solution exists when  $\mathbf{X}^T \mathbf{X}$  is invertible

$$\hat{\boldsymbol{\beta}}_{\text{OLS}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$



## Feature selection as motivation

---

Analytical solution exists when  $\mathbf{X}^T \mathbf{X}$  is invertible

$$\hat{\beta}_{\text{OLS}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$

The solution can be unstable or impossible to compute if

- ▶ there is **high correlation** between predictors, or
- ▶ if  $p > n$ .



## Feature selection as motivation

---

Analytical solution exists when  $\mathbf{X}^T \mathbf{X}$  is invertible

$$\hat{\beta}_{\text{OLS}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$$

The solution can be unstable or impossible to compute if

- ▶ there is **high correlation** between predictors, or
- ▶ if  $p > n$ .

**Solutions:** **Regularisation** or **feature selection**



## Filtering for feature selection

---

- ▶ Choose features through pre-processing
  - ▶ Features with maximum variance
  - ▶ Use only the first  $k$  PCA components



## Filtering for feature selection

---

- ▶ Choose features through pre-processing
  - ▶ Features with maximum variance
  - ▶ Use only the first  $k$  PCA components
- ▶ Examples of other useful measures
  - ▶ Use a univariate criterion, e.g. **F-score**: Features that correlate most with the response
  - ▶ **Mutual Information**: Reduction in uncertainty about  $x$  after observing  $y$
  - ▶ **Variable importance**: Determine variable importance with random forests



## Filtering for feature selection

- ▶ Choose features through pre-processing
  - ▶ Features with maximum variance
  - ▶ Use only the first  $k$  PCA components
- ▶ Examples of other useful measures
  - ▶ Use a univariate criterion, e.g. **F-score**: Features that correlate most with the response
  - ▶ **Mutual Information**: Reduction in uncertainty about  $x$  after observing  $y$
  - ▶ **Variable importance**: Determine variable importance with random forests
- ▶ **Summary**
  - ▶ **Pro**: Fast and easy
  - ▶ **Con**: Filtering mostly operates on single features and is not geared towards a certain method
  - ▶ Care with cross-validation and multiple testing necessary



## Filtering for feature selection

- ▶ Choose features through pre-processing
  - ▶ Features with maximum variance
  - ▶ Use only the first  $k$  PCA components
- ▶ Examples of other useful measures
  - ▶ Use a univariate criterion, e.g. **F-score**: Features that correlate most with the response
  - ▶ **Mutual Information**: Reduction in uncertainty about  $x$  after observing  $y$
  - ▶ **Variable importance**: Determine variable importance with random forests
- ▶ **Summary**
  - ▶ **Pro**: Fast and easy
  - ▶ **Con**: Filtering mostly operates on single features and is not geared towards a certain method
  - ▶ Care with cross-validation and multiple testing necessary
- ▶ Filtering is often more of a pre-processing step and less of a proper feature selection step



## Wrapping for feature selection

---

- ▶ **Idea:** Determine the best set of features by fitting models of different complexity and comparing their performance



## Wrapping for feature selection

---

- ▶ **Idea:** Determine the best set of features by fitting models of different complexity and comparing their performance
- ▶ **Best subset selection:** Try all possible (**exponentially many**) subsets of features and compare model performance with e.g. cross-validation



## Wrapping for feature selection

---

- ▶ **Idea:** Determine the best set of features by fitting models of different complexity and comparing their performance
- ▶ **Best subset selection:** Try all possible (**exponentially many**) subsets of features and compare model performance with e.g. cross-validation
- ▶ **Forward selection:** Start with just an intercept and add in each step the variable that improves fit the most (**greedy algorithm**)



## Wrapping for feature selection

---

- ▶ **Idea:** Determine the best set of features by fitting models of different complexity and comparing their performance
- ▶ **Best subset selection:** Try all possible (**exponentially many**) subsets of features and compare model performance with e.g. cross-validation
- ▶ **Forward selection:** Start with just an intercept and add in each step the variable that improves fit the most (**greedy algorithm**)
- ▶ **Backward selection:** Start with all variables included and then remove sequentially the one with the least impact (**greedy algorithm**)



## Wrapping for feature selection

---

- ▶ **Idea:** Determine the best set of features by fitting models of different complexity and comparing their performance
- ▶ **Best subset selection:** Try all possible (**exponentially many**) subsets of features and compare model performance with e.g. cross-validation
- ▶ **Forward selection:** Start with just an intercept and add in each step the variable that improves fit the most (**greedy algorithm**)
- ▶ **Backward selection:** Start with all variables included and then remove sequentially the one with the least impact (**greedy algorithm**)
- ▶ As discreet procedures, all of these methods **exhibit high variance** (small changes could lead to different predictors being selected, resulting in a potentially very different model)



## Embedding for feature selection

---

- ▶ **Embed/include** the feature selection into the model estimation procedure



## Embedding for feature selection

- ▶ **Embed/include** the feature selection into the model estimation procedure
- ▶ Ideally, penalization on the number of included features

$$\hat{\beta} = \arg \min_{\beta} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \sum_{j=1}^p \mathbb{1}(\beta_j \neq 0)$$

However, **discrete optimization problems** are hard to solve



## Embedding for feature selection

- ▶ **Embed/include** the feature selection into the model estimation procedure
- ▶ Ideally, penalization on the number of included features

$$\hat{\beta} = \arg \min_{\beta} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \sum_{j=1}^p \mathbb{1}(\beta_j \neq 0)$$

However, **discrete optimization problems** are hard to solve

- ▶ **Softer regularisation methods** can help

$$\hat{\beta} = \arg \min_{\beta} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \|\beta\|_q^q$$

where  $\lambda$  is a tuning parameter and  $q \geq 1$  or  $q = \infty$ .



# Feature selection

---

**Feature selection** can be addressed in multiple ways

- ▶ **Filtering:** Remove variables before the actual model for the data is built
  - ▶ Often crude but fast
  - ▶ Typically only pays attention to one or two features at a time (e.g. F-Score, MIC) or does not take the outcome variable into consideration (e.g. PCA)
- ▶ **Wrapping:** Consider the selected features as an additional hyper-parameter
  - ▶ computationally very heavy
  - ▶ most approximations are greedy algorithms
- ▶ **Embedding:** Include feature selection into parameter estimation through penalisation of the model coefficients
  - ▶ Naive form is equally computationally heavy as wrapping
  - ▶ **Soft-constraints** create biased but useful approximations



# Regularised regression

---



# Constrained and regularised regression

---

The optimization problem

$$\arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 \quad \text{subject to} \quad \|\boldsymbol{\beta}\|_q^q \leq t$$

for  $q > 0$  is equivalent to

$$\hat{\boldsymbol{\beta}} = \arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_q^q$$

when  $q \geq 1$ .



## Constrained and regularised regression

The optimization problem

$$\arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 \quad \text{subject to} \quad \|\boldsymbol{\beta}\|_q^q \leq t$$

for  $q > 0$  is equivalent to

$$\hat{\boldsymbol{\beta}} = \arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_q^q$$

when  $q \geq 1$ . This is the **Lagrangian** of the constrained problem.

**Note:** Constraints are convex for all  $q \geq 1$  but not differentiable in  $\boldsymbol{\beta} = \mathbf{0}$  for  $q = 1$ .



## Ridge regression

---

For  $q = 2$  the constrained problem is **ridge regression** (Tikhonov regularisation)

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = \arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_2^2$$

where  $\|\boldsymbol{\beta}\|_2^2 = \sum_{j=1}^p \beta_j^2$ .



## Ridge regression

For  $q = 2$  the constrained problem is **ridge regression** (Tikhonov regularisation)

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = \arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_2^2$$

where  $\|\boldsymbol{\beta}\|_2^2 = \sum_{j=1}^p \beta_j^2$ .

An **analytical solution** exists if  $\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p$  is invertible

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top \mathbf{y}$$



## Ridge regression

For  $q = 2$  the constrained problem is **ridge regression** (Tikhonov regularisation)

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = \arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_2^2$$

where  $\|\boldsymbol{\beta}\|_2^2 = \sum_{j=1}^p \beta_j^2$ .

An **analytical solution** exists if  $\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I}_p$  is invertible

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = (\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^T \mathbf{y}$$

If  $\mathbf{X}^T \mathbf{X} = \mathbf{I}_p$ , then

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = \frac{\hat{\boldsymbol{\beta}}_{\text{OLS}}}{1 + \lambda},$$

i.e.  $\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda)$  is **biased** but has **lower variance**.



## SVD and ridge regression

---

**Recall:** The SVD of a matrix  $\mathbf{X} \in \mathbb{R}^{n \times p}$  was

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$



## SVD and ridge regression

---

**Recall:** The SVD of a matrix  $\mathbf{X} \in \mathbb{R}^{n \times p}$  was

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$$

The analytical solution for ridge regression becomes ( $n \geq p$ )

$$\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) = (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top \mathbf{y}$$



## SVD and ridge regression

---

**Recall:** The SVD of a matrix  $\mathbf{X} \in \mathbb{R}^{n \times p}$  was

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$$

The analytical solution for ridge regression becomes ( $n \geq p$ )

$$\begin{aligned}\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) &= (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top \mathbf{y} \\ &= (\mathbf{V}\mathbf{D}^2\mathbf{V}^\top + \lambda \mathbf{I}_p)^{-1} \mathbf{V}\mathbf{D}\mathbf{U}^\top \mathbf{y}\end{aligned}$$



## SVD and ridge regression

---

**Recall:** The SVD of a matrix  $\mathbf{X} \in \mathbb{R}^{n \times p}$  was

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$$

The analytical solution for ridge regression becomes ( $n \geq p$ )

$$\begin{aligned}\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) &= (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top \mathbf{y} \\ &= (\mathbf{V}\mathbf{D}^2\mathbf{V}^\top + \lambda \mathbf{I}_p)^{-1} \mathbf{V}\mathbf{D}\mathbf{U}^\top \mathbf{y} \\ &= \mathbf{V}(\mathbf{D}^2 + \lambda \mathbf{I}_p)^{-1} \mathbf{D}\mathbf{U}^\top \mathbf{y}\end{aligned}$$



## SVD and ridge regression

**Recall:** The SVD of a matrix  $\mathbf{X} \in \mathbb{R}^{n \times p}$  was

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$$

The analytical solution for ridge regression becomes ( $n \geq p$ )

$$\begin{aligned}\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) &= (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top \mathbf{y} \\ &= (\mathbf{V}\mathbf{D}^2\mathbf{V}^\top + \lambda \mathbf{I}_p)^{-1} \mathbf{V}\mathbf{D}\mathbf{U}^\top \mathbf{y} \\ &= \mathbf{V}(\mathbf{D}^2 + \lambda \mathbf{I}_p)^{-1} \mathbf{D}\mathbf{U}^\top \mathbf{y} \\ &= \sum_{j=1}^p \frac{d_j}{d_j^2 + \lambda} \mathbf{v}_j \mathbf{u}_j^\top \mathbf{y}\end{aligned}$$



## SVD and ridge regression

**Recall:** The SVD of a matrix  $\mathbf{X} \in \mathbb{R}^{n \times p}$  was

$$\mathbf{X} = \mathbf{U}\mathbf{D}\mathbf{V}^\top$$

The analytical solution for ridge regression becomes ( $n \geq p$ )

$$\begin{aligned}\hat{\boldsymbol{\beta}}_{\text{ridge}}(\lambda) &= (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top \mathbf{y} \\ &= (\mathbf{V}\mathbf{D}^2\mathbf{V}^\top + \lambda \mathbf{I}_p)^{-1} \mathbf{V}\mathbf{D}\mathbf{U}^\top \mathbf{y} \\ &= \mathbf{V}(\mathbf{D}^2 + \lambda \mathbf{I}_p)^{-1} \mathbf{D}\mathbf{U}^\top \mathbf{y} \\ &= \sum_{j=1}^p \frac{d_j}{d_j^2 + \lambda} \mathbf{v}_j \mathbf{u}_j^\top \mathbf{y}\end{aligned}$$

Ridge regression **acts strongest** on principal components with **lower eigenvalues**, e.g. in presence of correlation between features.



## Effective degrees of freedom

---

Recall the **hat matrix**  $\mathbf{H} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top$  in OLS. The trace of  $\mathbf{H}$

$$\text{tr}(\mathbf{H}) = \text{tr}(\mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top) = \text{tr}(\mathbf{X}^\top \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1}) = \text{tr}(\mathbf{I}_p) = p$$

is equal to the trace of  $\hat{\Sigma}$  and the **degrees of freedom** for the regression coefficients.



## Effective degrees of freedom

Recall the **hat matrix**  $\mathbf{H} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top$  in OLS. The trace of  $\mathbf{H}$

$$\text{tr}(\mathbf{H}) = \text{tr}(\mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top) = \text{tr}(\mathbf{X}^\top \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1}) = \text{tr}(\mathbf{I}_p) = p$$

is equal to the trace of  $\hat{\Sigma}$  and the **degrees of freedom** for the regression coefficients.

In analogy define for ridge regression

$$\mathbf{H}(\lambda) := \mathbf{X}(\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top$$



## Effective degrees of freedom

Recall the **hat matrix**  $\mathbf{H} = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top$  in OLS. The trace of  $\mathbf{H}$

$$\text{tr}(\mathbf{H}) = \text{tr}(\mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top) = \text{tr}(\mathbf{X}^\top \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1}) = \text{tr}(\mathbf{I}_p) = p$$

is equal to the trace of  $\hat{\Sigma}$  and the **degrees of freedom** for the regression coefficients.

In analogy define for ridge regression

$$\mathbf{H}(\lambda) := \mathbf{X}(\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^\top$$

and

$$\text{df}(\lambda) := \text{tr}(\mathbf{H}(\lambda)) = \sum_{j=1}^p \frac{d_j^2}{d_j^2 + \lambda},$$

the **effective degrees of freedom**.



For  $q = 1$  the constrained problem is known as the **lasso**

$$\hat{\boldsymbol{\beta}}_{\text{lasso}}(\lambda) = \arg \min_{\boldsymbol{\beta}} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_1$$

- ▶ Smallest  $q$  in penalty such that constraint is still convex
- ▶ Produces **sparse solutions** (many coefficients exactly equal to zero) and therefore performs **feature selection**



## Intuition for the penalties (I)

Assume the OLS solution  $\beta_{\text{OLS}}$  exists and set

$$\mathbf{r} = \mathbf{y} - \mathbf{X}\beta_{\text{OLS}}$$

it follows for the **residual sum of squares (RSS)** that

$$\begin{aligned}\|\mathbf{y} - \mathbf{X}\beta\|_2^2 &= \|(\mathbf{X}\beta_{\text{OLS}} + \mathbf{r}) - \mathbf{X}\beta\|_2^2 \\ &= \|(\mathbf{X}(\beta - \beta_{\text{OLS}}) - \mathbf{r})\|_2^2 \\ &= (\beta - \beta_{\text{OLS}})^\top \mathbf{X}^\top \mathbf{X} (\beta - \beta_{\text{OLS}}) - 2\mathbf{r}^\top \mathbf{X} (\beta - \beta_{\text{OLS}}) + \mathbf{r}^\top \mathbf{r}\end{aligned}$$


which is an **ellipse** (at least in 2D) centred on  $\beta_{\text{OLS}}$ .




## Intuition for the penalties (II)

The least squares RSS is minimized for  $\beta_{OLS}$ . If a constraint is added ( $\|\beta\|_q^q \leq t$ ) then the RSS is minimized by the closest  $\beta$  possible that fulfills the constraint.

Lasso



Ridge



The blue lines are the contour lines for the RSS.







## Intuition for the penalties (III)

Depending on  $q$  the different constraints lead to different solutions. If  $\beta_{OLS}$  is in one of the coloured areas or on a line, the constrained solution will be at the corresponding dot.

**Sparsity** only for  $q \leq 1$

**Convexity** only for  $q \geq 1$





## Computational aspects of the Lasso (I)

---

What estimates does the lasso produce?



# Computational aspects of the Lasso (I)

---

What estimates does the lasso produce?

**Target function**

$$\arg \min_{\boldsymbol{\beta}} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_1$$



# Computational aspects of the Lasso (I)

What estimates does the lasso produce?

**Target function**

$$\arg \min_{\boldsymbol{\beta}} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_1$$

**Special case:**  $\mathbf{X}^\top \mathbf{X} = \mathbf{I}_p$ . Then

$$\frac{1}{2} \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 + \lambda \|\boldsymbol{\beta}\|_1 = \frac{1}{2} \mathbf{y}^\top \mathbf{y} - \underbrace{\mathbf{y}^\top \mathbf{X}}_{=\boldsymbol{\beta}_{\text{OLS}}^\top} \boldsymbol{\beta} + \frac{1}{2} \boldsymbol{\beta}^\top \boldsymbol{\beta} + \lambda \|\boldsymbol{\beta}\|_1 = g(\boldsymbol{\beta})$$



# Computational aspects of the Lasso (I)

What estimates does the lasso produce?

**Target function**

$$\arg \min_{\beta} \frac{1}{2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \|\beta\|_1$$

**Special case:**  $\mathbf{X}^\top \mathbf{X} = \mathbf{I}_p$ . Then

$$\frac{1}{2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \|\beta\|_1 = \frac{1}{2} \mathbf{y}^\top \mathbf{y} - \underbrace{\mathbf{y}^\top \mathbf{X}}_{=\beta_{\text{OLS}}^\top} \beta + \frac{1}{2} \beta^\top \beta + \lambda \|\beta\|_1 = g(\beta)$$

How do we find the solution  $\hat{\beta}$  in presence of the **non-differentiable** penalisation  $\|\beta\|_1$ ?



## Computational aspects of the Lasso (II)

---

For  $\mathbf{X}^\top \mathbf{X} = \mathbf{I}_p$  the target function can be written as

$$\arg \min_{\boldsymbol{\beta}} \sum_{j=1}^p -\beta_{\text{OLS},j} \beta_j + \frac{1}{2} \beta_j^2 + \lambda |\beta_j|$$



## Computational aspects of the Lasso (II)

For  $\mathbf{X}^T \mathbf{X} = \mathbf{I}_p$  the target function can be written as

$$\arg \min_{\boldsymbol{\beta}} \sum_{j=1}^p -\beta_{\text{OLS},j} \beta_j + \frac{1}{2} \beta_j^2 + \lambda |\beta_j|$$

This results in  $p$  **uncoupled** optimization problems.



## Computational aspects of the Lasso (II)

For  $\mathbf{X}^T \mathbf{X} = \mathbf{I}_p$  the target function can be written as

$$\arg \min_{\boldsymbol{\beta}} \sum_{j=1}^p -\beta_{\text{OLS},j} \beta_j + \frac{1}{2} \beta_j^2 + \lambda |\beta_j|$$

This results in  $p$  **uncoupled** optimization problems.

- ▶ **If**  $\beta_{\text{OLS},j} > 0$ , then  $\beta_j > 0$  to minimize the target
- ▶ **If**  $\beta_{\text{OLS},j} \leq 0$ , then  $\beta_j \leq 0$



## Computational aspects of the Lasso (II)

For  $\mathbf{X}^T \mathbf{X} = \mathbf{I}_p$  the target function can be written as

$$\arg \min_{\boldsymbol{\beta}} \sum_{j=1}^p -\beta_{\text{OLS},j} \beta_j + \frac{1}{2} \beta_j^2 + \lambda |\beta_j|$$

This results in  **$p$  uncoupled** optimization problems.

- ▶ **If**  $\beta_{\text{OLS},j} > 0$ , then  $\beta_j > 0$  to minimize the target
- ▶ **If**  $\beta_{\text{OLS},j} \leq 0$ , then  $\beta_j \leq 0$

Each case results in

$$\hat{\beta}_{\text{lasso},j} = \text{sign}(\beta_{\text{OLS},j})(|\beta_{\text{OLS},j}| - \lambda)_+ = \text{ST}(\beta_{\text{OLS},j}, \lambda),$$

where


- ▶  $x_+ = x$  if  $x > 0$  or 0 otherwise,
- ▶ and ST is called the **soft-thresholding operator**



## Relation to OLS estimates

Both ridge regression and the lasso estimates can be written as functions of  $\beta_{\text{OLS}}$  if  $\mathbf{X}^\top \mathbf{X} = \mathbf{I}_p$ .

$$\beta_{\text{ridge},j} = \frac{\beta_{\text{OLS},j}}{1 + \lambda} \quad \text{and} \quad \hat{\beta}_{\text{lasso},j} = \text{sign}(\beta_{\text{OLS},j})(|\beta_{\text{OLS},j}| - \lambda)_+$$



Visualisation of the transformations applied to the OLS estimates.



## Shrinkage and effective degrees of freedom

When  $\lambda$  is fixed, the **shrinkage** of the lasso estimate  $\beta_{\text{lasso}}(\lambda)$  compared to the OLS estimate  $\beta_{\text{OLS}}$  is defined as

$$s(\lambda) = \frac{\|\beta_{\text{lasso}}(\lambda)\|_1}{\|\beta_{\text{OLS}}\|_1}$$

**Note:**  $s(\lambda) \in [0, 1]$  with  $s(\lambda) \rightarrow 0$  for increasing  $\lambda$  and  $s(\lambda) = 1$  if  $\lambda = 0$



## Shrinkage and effective degrees of freedom

When  $\lambda$  is fixed, the **shrinkage** of the lasso estimate  $\beta_{\text{lasso}}(\lambda)$  compared to the OLS estimate  $\beta_{\text{OLS}}$  is defined as

$$s(\lambda) = \frac{\|\beta_{\text{lasso}}(\lambda)\|_1}{\|\beta_{\text{OLS}}\|_1}$$

**Note:**  $s(\lambda) \in [0, 1]$  with  $s(\lambda) \rightarrow 0$  for increasing  $\lambda$  and  $s(\lambda) = 1$  if  $\lambda = 0$

**Recall:** For ridge regression define

$$\mathbf{H}(\lambda) := \mathbf{X}(\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I}_p)^{-1} \mathbf{X}^T$$

and

$$\text{df}(\lambda) := \text{tr}(\mathbf{H}(\lambda)) = \sum_{j=1}^p \frac{d_j^2}{d_j^2 + \lambda},$$

the **effective degrees of freedom**.



## Prostate cancer dataset





Data to examine the correlation between the level of a prostate cancer-specific substance and a number of clinical measurements in men who just before partial or full removal of the prostate in patients.

- ▶  $n = 67$  samples
- ▶ A continuous response on the log-scale
- ▶  $p = 8$  features
  - ▶ e.g. log cancer volume, log prostate weight or age of patient



# Regularisation paths for varying $\lambda$

Red dashed lines indicate the  $\lambda$  selected by cross-validation





- ▶ In the general case, i.e.  $\mathbf{X}^T \mathbf{X} \neq \mathbf{I}_p$ , there is no explicit solution.



- ▶ In the general case, i.e.  $\mathbf{X}^T \mathbf{X} \neq \mathbf{I}_p$ , there is no explicit solution.
- ▶ Numerical solution possible, e.g. with **coordinate descent** where each  $\beta_j$  is updated separately with the remaining  $\beta_i$  with  $i \neq j$  fixed



- ▶ In the general case, i.e.  $\mathbf{X}^T \mathbf{X} \neq \mathbf{I}_p$ , there is no explicit solution.
- ▶ Numerical solution possible, e.g. with **coordinate descent** where each  $\beta_j$  is updated separately with the remaining  $\beta_i$  with  $i \neq j$  fixed
- ▶ As for ridge regression, **estimates are biased**



- ▶ In the general case, i.e.  $\mathbf{X}^T \mathbf{X} \neq \mathbf{I}_p$ , there is no explicit solution.
- ▶ Numerical solution possible, e.g. with **coordinate descent** where each  $\beta_j$  is updated separately with the remaining  $\beta_i$  with  $i \neq j$  fixed
- ▶ As for ridge regression, **estimates are biased**
- ▶ **Degrees of freedom** are equal to the number of non-zero coefficients



## Potential caveats of the lasso (I)

---

- ▶ **Sparsity of the true model:**

- ▶ The lasso only works if the data is generated from a sparse process.
- ▶ However, a dense process with many variables and not enough data or high correlation between predictors can be unidentifiable either way



## Potential caveats of the lasso (I)

---

- ▶ **Sparsity of the true model:**
  - ▶ The lasso only works if the data is generated from a sparse process.
  - ▶ However, a dense process with many variables and not enough data or high correlation between predictors can be unidentifiable either way
- ▶ **Correlations:** Many non-relevant variables correlated with relevant variables can lead to the selection of the wrong model, even for large  $n$



## Potential caveats of the lasso (I)

- ▶ **Sparsity of the true model:**

- ▶ The lasso only works if the data is generated from a sparse process.
- ▶ However, a dense process with many variables and not enough data or high correlation between predictors can be unidentifiable either way

- ▶ **Correlations:** Many non-relevant variables correlated with relevant variables can lead to the selection of the wrong model, even for large  $n$

- ▶ **Irrepresentable condition:** Split  $\mathbf{X}$  such that  $\mathbf{X}_1$  contains all **relevant variables** and  $\mathbf{X}_2$  contains all **irrelevant variables**. If

$$|(\mathbf{X}_2^T \mathbf{X}_1)(\mathbf{X}_1^T \mathbf{X}_1)^{-1}| < 1 - \eta$$

for some  $\eta > 0$  then the lasso is (almost) guaranteed to pick the true model



## Potential caveats of the lasso (II)

---

In practice, both the **sparsity of the true model** and the **irrepresentable condition** cannot be checked.

- ▶ Assumptions and domain knowledge have to be used



## Take-home message

---

- ▶ Filtering and wrapping methods useful for feature selection in practice but can be unprincipled or have high variance
- ▶ Regularised regression can help in numerically unstable situations (such as in ridge regression)
- ▶ The lasso can in addition perform variable selection