

Table of Laplace Transforms and trigonomerty

| $f(t)$ | $F(s)$ |
|-------------------------------------------|------------------------------------------------|
| $af(t) + bg(t)$ | $aF(s) + bG(s)$ |
| $tf(t)$ | $-F'(s)$ |
| $t^n f(t)$ | $(-1)^n F^{(n)}(s)$ |
| $e^{-at} f(t)$ | $F(s+a)$ |
| $f(t-T)\theta(t-T)$ | $e^{-Ts} F(s)$ |
| $f'(t)$ | $sF(s) - f(0)$ |
| $f''(t)$ | $s^2 F(s) - sf(0) - f'(0)$ |
| $f^{(n)}(t)$ | $s^n F(s) - \sum_{k=1}^n s^{n-k} f^{(k-1)}(0)$ |
| $\int_0^t f(\tau) d\tau$ | $\frac{F(s)}{s}$ |
| $\theta(t)$ | $\frac{1}{s}$ |
| $\frac{t^n}{n!}$ | $\frac{1}{s^{n+1}}$ |
| e^{-at} | $\frac{1}{s+a}$ |
| $\cosh(at)$ | $\frac{s}{s^2 - a^2}$ |
| $\sinh(at)$ | $\frac{a}{s^2 - a^2}$ |
| $\cos(bt)$ | $\frac{s}{s^2 + b^2}$ |
| $\sin(bt)$ | $\frac{b}{s^2 + b^2}$ |
| $\frac{t}{2b} \sin(bt)$ | $\frac{s}{(s^2 + b^2)^2}$ |
| $\frac{1}{2b^3} (\sin(bt) - bt \cos(bt))$ | $\frac{1}{(s^2 + b^2)^2}$ |

$$2\sin(a)\sin(b) = \cos(a-b) - \cos(a+b)$$

$$2\sin(a)\cos(b) = \sin(a-b) + \sin(a+b)$$

$$2\cos(a)\cos(b) = \cos(a-b) + \cos(a+b)$$