

# MVE550 2022 Lecture 1

## Introduction to stochastic processes

### Course introduction





Petter Mostad

Chalmers University

November 1, 2022

- ▶ Stochastic processes
- ▶ Bayesian inference
- ▶ Course structure and course content
- ▶ Dobrow Appendices A, B, C, D
- ▶ Dowbrow Chapter 1:
  - ▶ Conditional probability
  - ▶ Conditional expectation

# Things one might want to study



## Some common features in the examples

- ▶ There is a time involved. Observations “indexed” with a specific time.
- ▶ Possible *goals*: “Understand” something or *make predictions*.
- ▶ My opinion: Prediction is the central goal!
  - ▶ To “understand” something usually means to create some kind of underlying *model*.
  - ▶ Any model is a scientific model only if it makes *predictions*, and it can only be evaluated in terms of the correctness of its predictions.

# Deterministic and stochastic models

- ▶ Some models make exact predictions (without uncertainty).  
Example:  $F = ma$ .
- ▶ *Deterministic* models.
- ▶ In most cases, it is more reasonable to make probabilistic predictions.
- ▶ All examples above of this type.
- ▶ Stochastic models = probabilistic models, making probability predictions.

# Stochastic processes

- ▶ A stochastic process is a collection of *random variables*  $\{X_t, t \in I\}$ .
- ▶ The set  $I$  is the *index set* of the process.  $I$  most often represents a set of *specific times*.
- ▶ The random variables are defined on a common *state space*  $\mathcal{S}$ . This set represents the *possible values* the random variables  $X_t$  can have.
- ▶ In our four examples, the state spaces might be
  - ▶ A non-negative count.
  - ▶ A non-negative real number.
  - ▶ A set of species, with descriptions of their relevant genetic sequences and their relevant traits.
  - ▶ Some description of the amount of infections (and possibly immunity) in the population.
- ▶ Some further examples:
  - ▶ A vector of real numbers.
  - ▶ A grid of numbers (representing an image?)
  - ▶ A 3D grid of numbers (representing the stresses in a building?)
  - ▶ An infinite sequence of numbers.
  - ▶ A continuous function from  $[0, 1]$  to real numbers.

# The Markov property

- ▶ For us, the index set  $I$  will (generally) be some subset of the real numbers (representing time).
- ▶ Generally, for any  $t_0 \in I$ , the probabilities for outcomes for  $X_t$ , where  $t > t_0$ , may depend on the values of  $X_s$  for all  $s \leq t_0$ .
- ▶ The process fulfills the *Markov property* if, for any  $t_0 \in I$ , whenever  $X_{t_0}$  is known,  $X_t$  (with  $t > t_0$ ) is independent of the values for  $X_s$  for all  $s < t_0$ .
- ▶ More or less all the stochastic processes we will deal with in this course will have this property.

# What is a Random Variable?

Intuitive definition:

- ▶ A *variable* which has possible values in some *state space*  $\mathcal{S}$ . We will generally assume that the state space is a subset of the real numbers.
- ▶ Examples of state spaces used in the course:
  - ▶  $\mathcal{S} = \{1, 2, 3, 4\}$ .
  - ▶  $\mathcal{S}$  is all positive integers:  $\{1, 2, 3, 4, 5, \dots\}$ .
  - ▶  $\mathcal{S}$  is all non-negative real numbers.
- ▶ There are probabilities assigned to values and sets of values in the state space.
- ▶ We separate between *discrete* and *continuous* random variables.
- ▶ For discrete random variables, we assign a probability to each single value in the state space.
- ▶ For continuous random variables, we focus instead of assigning probabilities to *intervals* of values in the state space.
- ▶ (We will return shortly with more precise definitions.)

# Main types of stochastic processes in this course

Dobrow Chapters	Time ( $I$ )	State space ( $\mathcal{S}$ )
2&3: Discrete Markov chains	Discrete	Discrete
4: Branching processes	Discrete	Discrete
5: Markov chain Monte Carlo	Discrete	Continuous/Discrete
6: Poisson processes	Continuous	Discrete
7: Continuous-time Markov chains	Continuous	Discrete
8: Brownian motion	Continuous	Continuous

# What do we want to do with the models?

- ▶ Easiest approach: Set up model based on general knowledge, make predictions from models.
- ▶ Examples:
  - ▶ Throwing a dice.
  - ▶ Predictions about a card game.
  - ▶ Other types of game predictions.
- ▶ More useful situation:
  1. You have *data*.
  2. You want find a model so that the data could reasonably be produced by it.
  3. You want to use this model for predictions of future observations.
- ▶ **Using data in this way is called *inference*.**

# How to find a model that might have produced the data?

Two (main) alternatives:

- ▶ Classical inference (or “frequentist” inference):
  1. Find *estimates* for *parameters* of the model, using the data.
  2. To find *estimates*, use *estimators* that have *desireable properties*.
  3. Plug the estimates into the models and make predictions from resulting models.
- ▶ Bayesian inference:
  1. Set up a stochastic model making predictions of *observed data* and *possible future data*.
  2. Find the *conditional probability* for the future predictions given the values of the observed data.

- ▶ The Canvas pages!
- ▶ What is expected of you
- ▶ What you can expect from the course

- ▶ These appendices contain material that you (in principle) should know already.
- ▶ I *strongly recommend* that you look through these, at least to find out how much of them you know and how much and what you don't know.
- ▶ Appendix A: Getting started with R.
- ▶ Appendix B: Probability review.
- ▶ Appendix C: Summary of common probability distributions.
- ▶ Appendix D: Matrix algebra review.

A random variable  $X$  with state space  $S$  is a real-valued function on  $S$  together with a *probability*  $\Pr(\cdot)$  on  $S$ . The probability  $\Pr(\cdot)$  satisfies

- ▶  $0 \leq \Pr(A) \leq 1$  for all *measurable* subsets  $A \subseteq S$ .
- ▶  $\Pr(S) = 1$
- ▶  $\Pr(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \Pr(A_i)$  when the  $A_i$  are disjoint.
- ▶ These are the Kolmogorov axioms for probability.
- ▶ Measurable subsets are called *events*.
- ▶ What is a *measurable* subset?

# Measurable subsets

Let  $S$  be any set.

- ▶ A *sigma-algebra*  $\Omega$  on  $S$  is a set of subsets of  $S$  such that
  - ▶  $\Omega$  includes  $S$
  - ▶ If  $A \in \Omega$  then  $A^c = S \setminus A \in \Omega$ .
  - ▶ If  $A_1, A_2, \dots, \in \Omega$  then  $\cup_{i=1}^{\infty} A_i \in \Omega$
- ▶ The *measurable sets* are those that are in an appropriately defined sigma-algebra.
- ▶ What you need to know for this course: When  $S$  is finite or countable, all subsets will be measurable. When  $S$  is some interval of real numbers, there will exist subsets that are not measurable, but we will not be concerned with them.

- ▶ Note: Many random variables and stochastic processes can be represented with a computer program which *simulates* random output.
- ▶ The output is then *pseudo-random*
- ▶ We may then use

Frequency of computer output  $\approx$  Probability of output

- ▶ Making this precise yields powerful computational methods, some of which we will use and/or study in this course.

# Conditional probability

- ▶ Given events  $A$  and  $B$ , the *conditional probability* for  $A$  given  $B$  is

$$\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)}$$

- ▶ Events  $A$  and  $B$  are *independent* if  $\Pr(A \cap B) = \Pr(A) \Pr(B)$ .
- ▶ Law of total probability: Let  $B_1, \dots, B_k$  be a sequence of events that *partitions*  $S$ . Then

$$\Pr(A) = \sum_{i=1}^k \Pr(A \cap B_i) = \sum_{i=1}^k \Pr(A | B_i) \Pr(B_i).$$

- ▶ Bayes law for probabilities follows directly from definition above:

$$\Pr(B | A) = \frac{\Pr(A | B) \Pr(B)}{\Pr(A)}$$

# Notation for discrete probability distributions

- ▶ For a discrete random variable  $X$  we may write  $\Pr(X = x)$  for  $\Pr(\{x : X = x\})$ .
- ▶ For a joint distribution for two discrete random variables  $X$  and  $Y$  we may write  $\Pr(X = x, Y = y)$  for  $\Pr(\{x : X = x\} \cap \{y : Y = y\})$  and  $\Pr(X = x | Y = y)$  for  $\Pr(\{x : X = x\} | \{y : Y = y\})$
- ▶ The formulas of the previous overhead can then be written

$$\Pr(X = x | Y = y) = \frac{\Pr(X = x, Y = y)}{\Pr(Y = y)}$$

$$\Pr(X = x) = \sum_y \Pr(X = x | Y = y) \Pr(Y = y)$$

$$\Pr(Y = y | X = x) = \frac{\Pr(X = x | Y = y) \Pr(Y = y)}{\Pr(X = x)}$$

# The generic $\pi$ -notation

We may use the *generic*  $\pi$ -notation as a shorthand:

- ▶ Write  $\pi(x)$  for  $\Pr(X = x)$ ,  $\pi(x, y)$  for  $\Pr(X = x, Y = y)$  and  $\pi(x | y)$  for  $\Pr(X = x | Y = y)$ .
- ▶ The formulas of the previous overhead can then be written

$$\begin{aligned}\pi(x | y) &= \frac{\pi(x, y)}{\pi(y)} \\ \pi(x) &= \sum_y \pi(x | y)\pi(y) \\ \pi(y | x) &= \frac{\pi(x | y)\pi(y)}{\pi(x)}\end{aligned}$$

- ▶ The  $\pi(\cdot)$  notation will be used in the Lecture Notes, but is not used in Dobrow.

# Conditional densities for continuous distributions

- ▶ For a continuous random variable  $X$ , we will write its *density function* as  $\pi(x)$ , extending the generic  $\pi$  notation.
- ▶ If we have a joint distribution for continuous random variables  $X$  and  $Y$ , the joint density function may be written  $\pi(x, y)$ .
- ▶ We get formulas like

$$\int \pi(x) dx = 1 \quad \text{and} \quad \int \pi(x, y) dy = \pi(x).$$

- ▶ We may *define* the conditional density as

$$\pi(y | x) = \frac{\pi(x, y)}{\pi(x)}.$$

- ▶ We get similar formulas as for discrete variables:

$$\begin{aligned} \pi(x) &= \int_y \pi(x | y) \pi(y) dy \\ \pi(y | x) &= \frac{\pi(x | y) \pi(y)}{\pi(x)} \end{aligned}$$

# Expectation and conditional expectation

- ▶ Recall, the expectation of a discrete random variable is

$$E(Y) = \sum_y y\pi(y)$$

and of a continuous random variable

$$E(Y) = \int_y y\pi(y) dy.$$

- ▶ The conditional expectation in the discrete case is

$$E(Y | X = x) = \sum_y y\pi(y | x)$$

and in the continuous case

$$E(Y | X = x) = \int_y y\pi(y | x) dy.$$

# Law of total expectation

- ▶ If  $X$  is a discrete random variable, we get that

$$E(Y) = \sum_x E(Y | X = x) \pi(x).$$

- ▶ If  $X$  is a continuous random variable we get

$$E(Y) = \int_x E(Y | X = x) \pi(x) dx$$

- ▶ In both cases this can be written as

$$E(Y) = E(E(Y | X)).$$

# Law of total variance

- ▶ Recall that, by definition,

$$\text{Var}(Y) = E((Y - E(Y))^2) = E(Y^2) - E(Y)^2.$$

- ▶ Similarly, we have for the conditional variance

$$\text{Var}(Y | X = x) = E_{Y|X=x}((Y - E(Y | X = x))^2)$$

- ▶ With these definitions, we can now show the law of total variance:

$$\text{Var}(Y) = E(\text{Var}(Y | X)) + \text{Var}(E(Y | X))$$