

Matematisk Statistik och Diskret Matematik, MVE051/MSG810, VT20

Föreläsning 8

Nancy Abdallah

Chalmers - Göteborgs Universitet

Interval Estimate - Confidence Intervals

If we draw a random sample of size n from a population, then $\bar{x}$ is a point estimate of μ . It is more meaningful to estimate μ by an interval that we call confidence interval.

How to construct a Confidence Interval?

- Sampling from a normal distribution with known variance.
- Sampling from an unknown distribution (or from a non normal distribution).
- Sampling from a normal distribution with unknown variance.

Sampling from a normally distributed population

Suppose that the population is normally distributed with mean μ and variance σ^2 , and let $X_1, \dots, X_n$ be a random sample.

- The mean $\bar{X}$ is normally distributed with mean μ and variance σ^2/n .
- $\frac{\bar{X}-\mu}{\sigma/\sqrt{n}} \sim N(0, 1)$
- $P(-1.96 < \frac{\bar{X}-\mu}{\sigma/\sqrt{n}} < 1.96) = 0.95$ which is equivalent to $P(\bar{X} - 1.96 \frac{\sigma}{\sqrt{n}} < \mu < \bar{X} + 1.96 \frac{\sigma}{\sqrt{n}}) = 0.95$
- 95% of the intervals $(\bar{x} - 1.96 \frac{\sigma}{\sqrt{n}}, \bar{x} + 1.96 \frac{\sigma}{\sqrt{n}})$ contain the mean μ . This interval is called a 95% Confidence Interval (C.I.).

Example

Suppose a researcher, interested in obtaining an estimate of the average level of some enzyme in a certain human population, takes a sample of 10 individuals, determines the level of the enzyme in each, and computes a sample mean of $\bar{x} = 22$. Suppose further it is known that the variable of interest is approximately normally distributed with a variance of 45. Find a 95% interval confidence of μ .

Solution We Need to find the interval $(\bar{x} - 1.96\sigma/\sqrt{n}, \bar{x} + 1.96\sigma/\sqrt{n})$.
 $\sigma/\sqrt{n} = \sqrt{45/10} = 2.1213$, then the interval is

$$(22 - 1.96(2.1213), 22 + 1.96(2.1213))$$

which is

$$(17.84, 26.16)$$

Confidence interval of $\bar{x}$ from a normally distributed population with known σ

In general, a $100(1 - \alpha)$ confidence interval is given by

$$\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

where $z_{\alpha/2}$ is the value of z for which $P(z > z_{\alpha/2}) = \frac{\alpha}{2}$ in the standard normal distribution and is called the **the reliability coefficient**.

The C.I. can be written as

$$\text{estimator} \pm (\text{reliability coefficient}) \times (\text{standard error})$$

Interpretation

- **Probabilistic Interpretation:** In repeated sampling, from a normally distributed population with a known standard deviation, $100(1 - \alpha)$ percent of all intervals of the form $\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ will in the long run include the population mean μ .
- **Practical Interpretation:** When sampling is from a normally distributed population with known standard deviation, we are $100(1 - \alpha)$ percent confident that the single computed interval, $\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$, contains the population mean μ .

Example

A physical therapist wished to estimate, with 99 percent confidence, the mean maximal strength of a particular muscle in a certain group of individuals. He is willing to assume that strength scores are approximately normally distributed with a variance of 144. A sample of 15 subjects who participated in the experiment yield a mean of 84.3.

Solution $\alpha = 0.01$. We need to find $z_{\alpha/2}$, i.e. z such that $P(z > z_{\alpha/2}) = 0.005$. Using the table, we get $z_{\alpha/2} = 2.58$. Therefore, a 99% confidence interval is

$$(84.3 - (2.58)\sqrt{144/15}, 84.3 + (2.58)\sqrt{144/15})$$

which is

$$(76.3, 92.3)$$

Sampling from non-normal distribution

Usually the distribution of the population is not normal or is not even known. How do we construct a confidence interval in this case?

Definition (The central limit theorem)

Let $X_1, \dots, X_n$ be a random sample of size n from a distribution with mean μ and variance σ^2 . Then for large n , $\bar{X}$ is approximately normal with mean μ and variance σ^2/n .

Remark n is considered large if $n \geq 25$.

Example

Punctuality of patients in keeping appointments is of interest to a research team. In a study of patient flow through the offices of general practitioners, it was found that a sample of 35 patients was 17.2 minutes late for appointments, on the average. Previous research had shown the standard deviation to be about 8 minutes. The population distribution was felt to be nonnormal.

- 1 Approximate the distribution of the sample mean $\bar{X}$. What are the parameters?
- 2 What is the 90 percent confidence interval for μ , the true mean amount of time late for appointments?

Example (Solution)

- 1 Since $n \geq 25$ then $\bar{X}$ is approximately normally distributed with mean equals to the population mean μ (unknown) and standard deviation $\sigma/\sqrt{n} = 8/\sqrt{35} = 1.3522$
- 2 We need to find first $z_{\alpha/2} = z_{0.05}$. Using the table of standard normal distribution we get $z_{0.05}=1.645$. Hence, a 90% confidence interval is given by

$$(17.2 - 1.645(1.3522), 17.2 + 1.645(1.3522))$$

which is

$$(15.0, 19.4)$$

Distribution for population variance

Let $X_1, \dots, X_n$ be a random sample. Recall that $S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}$ is an unbiased estimate for the variance σ^2 .

Theorem

Suppose that $X_1, \dots, X_n$ are drawn from a normal distribution with mean μ and variance σ^2 . The random variable

$$(n-1)S^2/\sigma^2 = \sum_{i=1}^n (X_i - \bar{X})^2/\sigma^2$$

has a chi-squared distribution with $n-1$ degrees of freedom.

Confidence Interval for population variance

- To find a $100(1 - \alpha)\%$ C.I. for σ^2 , we find first a $100(1 - \alpha)\%$ C.I. for $(n - 1)S^2/\sigma^2$.
A C.I. for $(n - 1)S^2/\sigma^2$ is given by

$$(\chi_{1-\alpha/2}^2, \chi_{\alpha/2}^2)$$

i.e. $P(\chi_{1-\alpha/2}^2 \leq (n - 1)S^2/\sigma^2 \leq \chi_{\alpha/2}^2) = 1 - \alpha$

- By using operation on inequalities, we obtain the C.I on σ^2 .
A $100(1 - \alpha)\%$ C.I. for σ^2 is then

$$\left(\frac{(n - 1)S^2}{\chi_{\alpha/2}^2}, \frac{(n - 1)S^2}{\chi_{1-\alpha/2}^2} \right)$$

Example

Given the following data

9.7, 12.3, 11.2, 5.1, 24.8, 14.8, 17.7

Find a 95% confidence interval for the variance and for the standard deviation.

Solution The sample variance is $s^2 = 39.763$. The degrees of freedom are $n - 1 = 6$ and $\alpha = 0.05$. From the table, we get $\chi_{0.975}^2 = 1.237$ and $\chi_{0.025}^2 = 14.449$. Hence, our 95% C.I for σ^2 is

$$\left(\frac{6(39.763)}{14.449}, \frac{6(39.763)}{1.237} \right) = (16.512, 192.868)$$

and the 95% C.I. for σ is

$$(4.063, 13.888)$$

Student T-distribution

- Let Z and X be independent random variables such that $Z \in N(0, 1)$ and $X \in \chi^2_\gamma$. Then

$$T = \frac{Z}{\sqrt{X^2/\gamma}}$$

is said to have a Student T distribution with γ degrees of freedom.

- Properties of T-distribution:
 - The graph of the density function is a bell-shape symmetrical about the mean which is equal to 0.
 - It approaches the normal distribution as n increases.
 - The variance of the t distribution is greater than 1. If $df > 2$, the variance is equal to $\frac{df}{df-2}$ where df is the degrees of freedom.

Confidence interval of $\bar{x}$ from a normally distributed population with unknown σ

Let $X_1, \dots, X_n$ be a random sample drawn from a normally distributed population with unknown μ and σ .

- It is easy to show that $\frac{\bar{X}-\mu}{S/\sqrt{n}}$ follows a T distribution with $n-1$ degrees of freedom.
- To find a $100(1-\alpha)\%$ C.I., we use the same method as for the normal distribution, i.e.

$$\bar{x} \pm t_{(\alpha/2)} s / \sqrt{n}$$

where $\bar{x}$ is the mean of the sample, s its standard deviation, and $t_{(\alpha/2)}$ is the value of t for which $P(t > t_{(\alpha/2)}) = \alpha/2$.

- The values of t can be found using table VI p.699-700.

Example

A set of data on the sulfur dioxide concentration (in micrograms per cubic meter) in a Bavarian forest has been collected. The sample contains 24 values, its mean is $53.92 \mu\text{g}/\text{m}^3$, the sample variance is 101.480 and the standard deviation is $10.07 \mu\text{g}/\text{m}^3$. To obtain a 95% C.I. we find $t_{\alpha/2}$ from the table. $t_{0.025} = 2.069$. Hence, a 95% C.I. is given by

$$(53.92 - 2.069(10.07)/\sqrt{24}, 53.92 + 2.069(10.07)/\sqrt{24})$$

which is equal to

$$(49.67, 58.17)$$

Summary - C.I. for the population mean

